

SRB MEASURES FOR HYPERBOLIC POLYGONAL BILLIARDS

GIANLUIGI DEL MAGNO, JOÃO LOPES DIAS, PEDRO DUARTE,
JOSÉ PEDRO GAIVÃO, AND DIOGO PINHEIRO

ABSTRACT. We prove that polygonal billiards with contracting reflection laws exhibit hyperbolic attractors with countably many ergodic SRB measures. These measures are robust under small perturbations of the reflection law, and the tables for which they exist form a generic set in the space of all polygons. Specific polygonal tables are studied in detail.

1. INTRODUCTION

A planar billiard is the dynamical system describing the motion of a point particle moving freely in the interior of a connected compact subset $P \subset \mathbb{R}^2$ with piecewise smooth boundary. The particle slides along straight lines until it hits the boundary ∂P . If the collision occurs at a smooth boundary point, then the particle gets reflected according to a prescribed rule, called the reflection law. Otherwise, the particle stops and stays forever at the non-smooth boundary point where the collision occurred. The billiard dynamics is not smooth: the singularities of the dynamics are generated by non-smooth boundary points and trajectories having tangential collisions with ∂P . The reflection law most commonly considered is the specular one: the tangential component of the particle velocity remains the same, while the normal component changes its sign. Nevertheless, there exist alternative reflection laws that are equally reasonable and have interesting mathematical implications. In this paper, we study a family of these alternative laws.

One of the interesting features of billiards is that they exhibit a broad spectrum of dynamics. On one end of the spectrum, there are the integrable billiards. The only known integrable billiard tables are ellipses, rectangles, equilateral triangles, right isosceles triangles and right triangles with an angle $\pi/6$. Polygonal billiards are close to integrable systems in the sense that all their Lyapunov exponents are zero, which is a direct consequence of having zero topological entropy [17]. In particular, polygonal billiards are never hyperbolic. For some nice reviews on polygonal billiards, see [14, 29, 31]. On the other end of the spectrum, there are the hyperbolic billiards, i.e. billiards with non-zero

Lyapunov exponents. The Sinai billiards were the first examples of hyperbolic billiards; their tables consist of tori with a finite number of convex obstacles [26]. Another important class of hyperbolic billiards is represented by the semi-focusing billiards, so called because their tables are formed by convex (outward) curves and possibly straight segments. The most famous examples are the Bunimovich stadium and the Wojtkowski cardioid [7, 8, 33]. All the three examples of hyperbolic billiards mentioned above have strong ergodic properties; they are Bernoulli systems, and in particular ergodic and K-mixing. Other semi-focusing hyperbolic billiards were discovered by Markarian and Donnay [13, 22]. Examples of hyperbolic billiards of great interest in statistical mechanics are the Lorentz gases and hard-ball systems [28]. For a recent and rather complete account on the theory of hyperbolic billiards, we refer the reader to [11].

In this paper, we are interested in planar polygonal billiards with contracting reflection laws. This means that the function describing the dependence of the reflection angle on the incidence angle is a contraction. The simplest example of such a law is the linear contraction $\theta \mapsto \sigma\theta$ with $0 < \sigma < 1$, already studied in [4, 5, 15, 21]. To distinguish polygonal billiards with contracting reflection laws from those with specular reflection law, we will refer to the former as contracting polygonal billiards and to the latter as standard polygonal billiards.

The dynamics of these two classes of billiards are very different. Standard polygonal billiards exhibit the characteristic features of parabolic systems: zero entropy, subexponential growth of periodic orbits and subexponential divergence of nearby orbits. Instead, as we show in this paper, contracting polygonal billiards are typically uniformly hyperbolic systems (with singularities), and so they have positive entropy, exponential divergence of orbits, dense periodic orbits, and countably many ergodic and mixing components. Moreover, contracting polygonal billiards may exhibit mixed behavior, with hyperbolic and non-hyperbolic attractors coexisting. Yet another essential difference between these two classes of billiards is that standard polygonal billiards preserve the phase space area, whereas the contracting ones do not, this being due to the phase space contraction produced by the new reflection law. Therefore, an important part of the analysis of contracting polygonal billiards is represented by the study of their invariant measures. In this paper, we give sufficient conditions for the existence of Sinai-Ruelle-Bowen measures (SRB measures for short), which are measures with non-zero Lyapunov exponents and absolutely conditional measures on unstable manifolds. Our results rely on a general theory of SRB measures for uniformly systems with singularities developed by Pesin [23] and Sataev [25].

The first results on billiards with contracting reflection laws were obtained by Markarian, Pujals and Sambarino. In [21], they proved

that many billiards with contracting reflection law, including polygonal billiards, have dominated splitting, i.e. the tangent bundle of their phase space splits into two invariant directions such that the growth rate along one direction dominates uniformly the growth rate along the other direction. Note that uniform hyperbolicity implies dominated splitting. Markarian, Pujals and Sambarino also obtained some results concerning the hyperbolicity of polygonal billiards with strongly contracting reflection laws. In this paper, we improve their results. Specific polygonal tables and linear contracting reflection laws were studied in [5] and [15]. In the first of these two papers, the existence of a unique SRB measure was established for the equilateral triangle using a natural Markov partition. In the second one, we proved the existence of horseshoes and several properties of hyperbolic periodic orbits for the square. Works on billiards related to ours are the one of Zhang, where she proved the existence of SRB measures and obtained Green-Kubo formulae for two-dimensional periodic Lorentz gases with reflection laws modified according to certain ‘twisting’ rules [37], and the one of Chernov, Karepanov and Simányi on a gas of hard disks moving in a two-dimensional cylinder with non-elastic collisions between the disks and the walls [12]. Finally, see also [3, 4], for numerical results on billiards with non-specular reflections laws relevant in the study of certain optical systems.

We now describe the organization of this paper and at the same time our results in more detail. In Section 2, we give the definitions of the billiard map with general reflection law and of a contracting reflection law. We then discuss basic properties of the billiard map, including its singular sets, derivative and attractors.

The general hyperbolic properties of contracting polygonal billiards are discussed in Section 3. There, we prove that every contracting polygonal billiard has a dominated splitting, and provide conditions for the existence of hyperbolic sets. As a corollary, we obtain that if a polygon does not have parallel sides, then the correspondent contracting billiard has hyperbolic attractors.

Sufficient conditions for the existence of SRB measures on hyperbolic attractors are given in Section 4. As already mentioned, the result we obtain relies on general results of Pesin and Sataev. To be able to use these results, we show that contracting polygonal billiards satisfy a crucial condition, which roughly speaking, amounts to requiring that billiard trajectories do not visit too often a small neighborhood of the vertexes of the polygon. In the same section, we also prove that the existence of an SRB measure is a robust property under small perturbation of the reflection law.

In Section 5, we prove that polygonal billiards with strongly contracting reflection laws have generically countably many ergodic SRB

measures. We then obtain the same conclusion for every regular polygon with an odd number of sides in Section 6. To arrive to these results, we use a ‘perturbation argument’. Indeed, for strongly contracting reflection laws, the billiard map is close to the one-dimensional map obtained as the limit of the billiard map when the contraction becomes infinitely large. This map is called the slap map, and was introduced in [21]. If a polygon does not have parallel sides, then the slap map is expanding, and so the billiard map remains hyperbolic provided that the reflection law is strongly contracting. In Section 6, we also derive some general results about the absence of hyperbolic attractors when the billiard table is a regular polygon with an even number of sides.

Section 7 is devoted to the proof of the existence of SRB measures for acute triangles. The simpler dynamics of this class of polygons allows us to obtain an explicit estimate on the strength of the contraction needed for the existence of SRB measures. Finally, in Section 8, we give sufficient conditions for the existence and absence of hyperbolic attractors in rectangular billiards.

2. PRELIMINARIES

In this section, we define the mathematical objects we intend to study. First, we give a detailed construction of the billiard map for polygonal tables and general reflection laws. Then, we introduce a class of contracting reflection laws, and for such laws, we derive a series of preliminary results concerning the singular sets, the derivative and the attracting sets of polygonal billiard maps.

2.1. Billiard map. Let P be a non self-intersecting n -gon. Denote by $|\partial P|$ the length of ∂P and by C_1, \dots, C_n the vertexes of P . Let $\zeta: [0, |\partial P|] \rightarrow \mathbb{R}^2$ be the positively oriented parametrization of the curve ∂P by arclength such that if \tilde{s}_i is the arclength parameter corresponding to C_i , then $0 = \tilde{s}_1 < \dots < \tilde{s}_n < \tilde{s}_{n+1} = |\partial P|$. Note that the parametrization ζ is smooth everywhere on $[0, |\partial P|]$ except at $\tilde{s}_2, \dots, \tilde{s}_n$, and that $\zeta(0) = \zeta(|\partial P|)$, because ∂P is a closed curve.

Consider the following subsets of \mathbb{R}^2 ,

$$M = (0, |\partial P|) \times \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \quad \text{and} \quad V = \{\tilde{s}_1, \dots, \tilde{s}_{n+1}\} \times \left(-\frac{\pi}{2}, \frac{\pi}{2}\right).$$

For every $x = (s, \theta) \in M \setminus V$, define $q(x) = \zeta(s)$, and let $v(x)$ be the unit vector of \mathbb{R}^2 such that the oriented angle formed by $\zeta'(s)$ and $v(x)$ is equal to $\pi/2 - \theta$. Thus, $(q(x), v(x))$ defines a unit tangent vector of \mathbb{R}^2 pointing inside P . Both pairs (s, θ) and $(q(x), v(x))$ specify the state of the billiard particle immediately after a collision with ∂P . For this reason, we will refer to both (s, θ) and $(q(x), v(x))$ as the *collisions* of the billiard particle.

Given $x \in M \setminus V$ and $\tau > 0$, denote by $(q(x), q(x) + \tau v(x))$ the open segment of \mathbb{R}^2 with endpoints $q(x)$ and $q(x) + \tau v(x)$. Next, define

$$A(x) = \{\tau > 0: (q(x), q(x) + \tau v(x)) \subset P\}.$$

This set is nonempty and bounded, and so we can define

$$t(x) = \sup A(x) \quad \text{and} \quad q_1(x) = q(x) + t(x)v(x).$$

It is easy to see that $q_1(x)$ is the point where the particle leaving the point $q(x)$ with velocity $v(x)$ hits ∂P , whereas $t(x)$ is the euclidean distance between $q(x)$ and $q_1(x)$.

Let S_1^+ be the closure in \mathbb{R}^2 of all collisions mapped by q_1 into vertexes of P , i.e.,

$$S_1^+ = \bigcup_{i=1}^n \overline{q_1^{-1}(C_i)}.$$

Also, let

$$N_1^+ = \partial M \cup V \cup S_1^+.$$

For every $x \in M \setminus N_1^+$, define

$$s_1(x) = \zeta^{-1}(q_1(x))$$

and

$$\bar{v}_1(x) = -v(x) + 2\langle \zeta'(s_1(x)), v(x) \rangle \zeta'(s_1(x)),$$

where $\langle \cdot, \cdot \rangle$ is the euclidean scalar product of \mathbb{R}^2 . The billiard particle leaving $q(x)$ in the direction $v(x)$ hits ∂P at $q_1(x)$, and undergoes a reflection. When this reflection is specular, i.e., the reflection and the incidence angles coincide, the velocity of the particle immediately after the collision is equal to $\bar{v}_1(x)$. The angle formed by $v_1(x)$ with $\zeta'(s_1(x))$ is given

$$\bar{\theta}_1(x) = \arcsin(\langle \zeta'(s_1(x)), \bar{v}_1(x) \rangle).$$


Let L_0 and L_1 be the sides of P containing $q(x)$ and $q_1(x)$, respectively. Also, let ℓ_0 and ℓ_1 be the lines through the origin of \mathbb{R}^2 parallel to L_0 and L_1 , respectively. We assume that ℓ_0 and ℓ_1 have the same orientation of L_0 and L_1 . Denote by $\delta(L_0, L_1)$ the smallest positive angle of the rotation that maps ℓ_0 to ℓ_1 counterclockwise. We call such an angle the *angle formed by the sides L_0 and L_1* . It is not difficult to deduce that the relation between $\bar{\theta}_1(x)$ and $\theta(x)$ read as

$$\bar{\theta}_1(x) = \pi - \delta(L_0, L_1) - \theta(x). \quad (2.1)$$

The billiard map $\bar{\Phi}: M \setminus N_1^+ \rightarrow M$ with the specular reflection law is given by

$$\bar{\Phi}(x) = (s_1(x), \bar{\theta}_1(x)) \quad \text{for } x \in M \setminus N_1^+.$$

This map is a smooth embedding [11, Theorem 2.33].

FIGURE 1. Billiard map Φ_f .

2.2. General reflection laws. We now give the definition of the billiard map with general reflection law. A reflection law is given by a function $f: (-\pi/2, \pi/2) \rightarrow (-\pi/2, \pi/2)$. If we define $\theta_1(x) = f(\bar{\theta}_1(x))$ for $x \in M \setminus N_1^+$, then the billiard map $\Phi_f: M \setminus N_1^+ \rightarrow M$ with reflection law f (see Fig. 1) is given by

$$\Phi_f(x) = (s_1(x), \theta_1(x)) \quad \text{for } x \in M \setminus N_1^+.$$

Let $R_f(s, \theta) = (s, f(\theta))$ for every $(s, \theta) \in M$. So Φ_f may be written as

$$\Phi_f = R_f \circ \bar{\Phi}.$$

It follows that Φ_f is a C^k embedding with $k \geq 1$ provided that f is.

2.3. Contracting reflection laws. Given a differentiable function $f: (-\pi/2, \pi/2) \rightarrow \mathbb{R}$, let

$$\lambda(f) = \sup_{\theta \in (-\pi/2, \pi/2)} |f'(\theta)|.$$

Denote by \mathcal{R} the set of all differentiable $f: (-\pi/2, \pi/2) \rightarrow \mathbb{R}$ such that $f(0) = 0$ and $\lambda(f) < \infty$. For every transformation $\Psi: M \setminus N_1^+ \rightarrow \mathbb{R}^2$, define

$$\|\Psi\|_\infty = \sup_{x \in M \setminus N_1^+} \|\Psi(x)\|_2,$$

where $\|\cdot\|_2$ is the euclidean norm of \mathbb{R}^2 . Let X be the set of all $\Psi: M \setminus N_1^+ \rightarrow \mathbb{R}^2$ such that $\|\Psi\|_\infty < \infty$. The spaces (\mathcal{R}, λ) and $(X, \|\cdot\|_\infty)$ are normed space.

Consider the map from \mathcal{R} to X defined by $f \mapsto \Phi_f$. This map is continuous. Indeed, if f_1 and f_2 belong to \mathcal{R} , then

$$\begin{aligned} \|\Phi_{f_1} - \Phi_{f_2}\|_\infty &\leq \|R_{f_1} - R_{f_2}\|_\infty \\ &\leq \sup_{\theta \in (-\pi/2, \pi/2)} |f_1(\theta) - f_2(\theta)| \\ &\leq \frac{\pi}{2} \lambda(f_1 - f_2). \end{aligned}$$

For each $k \geq 1$, we define \mathcal{R}_1^k to be the set of all $f \in \mathcal{R}$ that are C^k embeddings with $\lambda(f) < 1$. If $f \in \mathcal{R}_1^k$, then f is a strict contraction, and $\theta = 0$ is its unique fixed point. Since f is monotone, it admits a




FIGURE 2. Contracting reflection law

continuous extension up to $-\pi/2$ and $\pi/2$, which is naturally denoted by $f(-\pi/2)$ and $f(\pi/2)$.

Strictly increasing reflection laws $f \in \mathcal{R}_1^k$ were also considered in [21]. The simplest example of such a law is $f(\theta) = \sigma\theta$ with $0 < \sigma < 1$ [4, 15, 21]. See Fig. 2.

From now on, we assume that $f \in \mathcal{R}_1^k$ with $k \geq 1$ unless otherwise stated.

2.4. Singular sets. A curve γ contained in M is called *strictly decreasing* (resp. *strictly increasing*), if γ is the graph of a continuous strictly decreasing (resp. *strictly increasing*) function $h: I \rightarrow (-\pi/2, \pi/2)$ with I being an interval of $(0, |\partial Q|)$.

In the next two propositions, we state the main geometrical properties of the sets S_1^+ and S_1^- . Their proofs are given in the Appendix.

Proposition 2.1. *The set S_1^+ is a union of finitely many analytic compact curves $\Gamma_1, \dots, \Gamma_m$ such that*

- (1) $\Gamma_i \subset [\tilde{s}_k, \tilde{s}_{k+1}] \times (-\pi/2, \pi/2)$ for some $1 \leq k \leq n$,
- (2) Γ_i is strictly decreasing,
- (3) Γ_i and Γ_j intersect transversally for $i \neq j$,
- (4) $\Gamma_i \cap \Gamma_j \subset \partial\Gamma_i \cup \partial\Gamma_j$ for $i \neq j$.

Let $F(s, \theta) = (s, f(-\theta))$ for $(s, \theta) \in M$, and define

$$S_1^- = F(S_1^+).$$

Since F is a C^k embedding, S_1^+ and S_1^- are diffeomorphic.

Proposition 2.2. *The conclusions of Proposition 2.1 hold for S_1^- if $f' < 0$, and hold with ‘decreasing’ replaced by ‘increasing’ in conclusion (3) if $f' > 0$.*

The iterates of the sets of S_1^+ and S_1^- play a crucial role in the study of the properties of the billiard map. Hence, for $n \geq 1$, define recursively

$$S_{n+1}^+ = S_n^+ \cup \Phi_f^{-1}(S_n^+) \quad \text{and} \quad S_{n+1}^- = S_n^- \cup \Phi_f(S_n^-).$$

Let $N_1^- = \partial(\Phi_f(M \setminus N_1^+))$. Note that while the set N_1^+ does not depend on the reflection law f , the set N_1^- does. For later use, we also

introduce the iterates of N_1^+ and N_1^- . For $n \geq 1$, set

$$N_{n+1}^+ = N_n^+ \cup \Phi_f^{-1}(N_n^+) \quad \text{and} \quad N_{n+1}^- = N_n^- \cup \Phi_f(N_n^-).$$

The set N_n^+ (resp. N_n^-) consist of the points of M where the map Φ_f^n (resp. Φ_f^{-n}) is not defined, and is called the *singular set* of Φ_f^n (resp. Φ_f^{-n}). Finally, let $N_\infty^+ = \bigcup_{n \geq 1} N_n^+$, and let $N_\infty^- = \bigcup_{n \geq 1} N_n^-$.

2.5. Derivative of the billiard map. Given $(s_0, \theta_0) \in M \setminus N_1^+$, let $(s_1, \theta_1) = \Phi_f(s_0, \theta_0)$. Recall that $t(s_0, \theta_0)$ denotes the length of the segment connecting $q(s_0, \theta_0)$ and $q_1(s_0, \theta_0)$. For the computation of the differential of $\bar{\Phi}$, see for instance [11, Formula (2.26)]. Since $D\Phi_f = DR_f \cdot D\bar{\Phi}$, we easily obtain

$$D\Phi_f(s_0, \theta_0) = - \begin{pmatrix} \frac{\cos \theta_0}{\cos \bar{\theta}_1} & \frac{t(s_0, \theta_0)}{\cos \bar{\theta}_1} \\ 0 & f'(\bar{\theta}_1) \end{pmatrix}.$$

Now, suppose that $(s_0, \theta_0) \in M \setminus N_n^+$ for some $n > 0$. Let $(s_i, \theta_i) = \Phi_f^i(s_0, \theta_0)$ for $i = 1, \dots, n$. We easily see that

$$D\Phi_f^n(s_0, \theta_0) = (-1)^n \begin{pmatrix} \alpha_n(s_0, \theta_0) & \gamma_n(s_0, \theta_0) \\ 0 & \beta_n(s_0, \theta_0) \end{pmatrix}, \quad (2.2)$$

where

$$\alpha_n(s_0, \theta_0) = \frac{\cos \theta_0}{\cos \bar{\theta}_n} \prod_{i=1}^{n-1} \frac{\cos \theta_i}{\cos \bar{\theta}_i}, \quad \beta_n(s_0, \theta_0) = \prod_{i=1}^n f'(\bar{\theta}_i),$$

and

$$\gamma_n(s_0, \theta_0) = \sum_{i=0}^{n-1} \beta_i(s_0, \theta_0) \frac{t(s_i, \theta_i)}{\cos \bar{\theta}_n} \prod_{k=i+1}^{n-1} \frac{\cos \theta_k}{\cos \bar{\theta}_k}.$$

Let $\rho: (-\pi/2, \pi, 2) \rightarrow [1, +\infty)$ be the function defined by

$$\rho(\theta) = \frac{\cos(f(\theta))}{\cos \theta} \quad \text{for } \theta \in (-\pi/2, \pi, 2).$$

Since $f \in \mathcal{R}_1^k$, it follows easily that ρ is continuous, and that $\rho(\theta) \geq 1$ with equality if and only if $\theta = 0$. Also, from $\lim_{\theta \rightarrow \pm\pi/2} \rho(\theta) = +\infty$, we obtain

$$r(\epsilon) := \min_{\epsilon \leq |\theta| < \pi/2} \rho(\theta) > 1 \quad \text{for } 0 < \epsilon < \pi/2.$$

Define

$$\Lambda_{n-1}(s_1, \theta_1) = \prod_{i=1}^{n-1} \rho(\bar{\theta}_i).$$

From the definition of α_n , we see that

$$\alpha_n(s_0, \theta_0) = \frac{\cos \theta_0}{\cos \bar{\theta}_n} \cdot \Lambda_{n-1}(s_1, \theta_1). \quad (2.3)$$

2.6. Attractors. Let

$$D_f = \bigcap_{n \geq 0} \Phi_f^n(M \setminus N_\infty^+).$$

It can be checked that $\Phi_f^{-1}(D_f) = D_f$, which in turn implies that $D_f \cap N_\infty^- = \emptyset$. Thus, every element of D_f has infinite positive and negative semi-orbits. We say that a subset Σ of M is *invariant* (resp. *forward invariant*) if $\Sigma \subset D_f$ and $\Phi_f^{-1}(\Sigma) = \Sigma$ (resp. $\Phi_f(\Sigma) \subset \Sigma$). Following the terminology introduced by Pesin in his work on maps with singularities [23], we call the closure of D_f the *attractor* of Φ_f .

Lemma 2.3. *We have*

$$D_f \subset \tilde{D}_f := \left\{ (s, \theta) \in M : |\theta| < \frac{\pi}{2} \lambda(f) \right\}.$$

Proof. The conclusion is a direct consequence of the invariance of D_f , and the fact that the absolute value of the angle of every element of $\Phi_f(D_f)$ is not larger than $\pi \lambda(f)/2$. \square

Definition 2.1. We say that two sides L_1 and L_2 of P *see each other* if there exists a line segment of a billiard trajectory whose endpoints lie in the interior of L_1 and L_2 .

Let ν be the Lebesgue measure $dsd\theta$ of \mathbb{R}^2 . When the reflection law is the standard one, i.e., $f = 1$, we have $D_f = M \setminus (N_\infty^- \cup N_\infty^+)$, and $\cos \theta dsd\theta$ is the natural invariant measure for Φ_f . In this case, it is easy to see that $\nu(D_f) = \nu(M) > 0$. For $f \in \mathcal{R}_1^k$, the situation is more delicate; we may have $\nu(D_f) = 0$, and therefore only singular invariant measures for Φ_f . The measures that play a prominent role in the study of hyperbolic systems are the SRB measures. We will prove the existence of these measures in Section 4. In the following propositions, we give sufficient conditions for $\nu(D_f) = 0$.

Proposition 2.4. *Suppose that P has the property*

$$|\pi - \delta(L_i, L_j)| \neq \pi/2$$

for all the pairs of sides L_i and L_j seeing each other. Then $\nu(D_f) = 0$ for $\lambda(f)$ sufficiently small.

Proposition 2.5. *Suppose that*

$$\sup_{\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})} \frac{|f'(\theta)|}{\cos \theta} < 1.$$

Then $\nu(D_f) = 0$.

Remark 2.6. An example of a reflection law that satisfies the hypothesis of Proposition 2.5 is $f(\theta) = \sigma \sin \theta$ with $0 < \sigma < 1$.

2.7. Notation. Instead of $\Phi_f, \lambda(f), D_f$, we write Φ, λ, D when no confusion can arise. We also write $x_n = (s_n, \theta_n)$ for $\Phi_f^n(x_0)$ with $x_0 = (s_0, \theta_0)$.

3. HYPERBOLICITY

Let $u \in T_x M$ with $x \in M$, and denote by u_s and u_θ the components of u with respect to the coordinates (s, θ) . By setting $\|u\|^2 = u_s^2 + u_\theta^2$ for every $u \in T_x M$, we introduce a differentiable norm $\|\cdot\|$ on M . Let d be the metric on M generated by $\|\cdot\|$.

We say that an invariant set $\Sigma \subset M$ has a *dominated splitting* if there exist a non-trivial continuous splitting $T_\Sigma M = E \oplus F$ and two constants $0 < \mu < 1$ and $A > 0$ such that for all $x \in \Sigma$ and $n \geq 1$, we have

$$\|D\Phi^n|_{E(x)}\| \cdot \|D\Phi^{-n}|_{F(\Phi^n x)}\| \leq A\mu^n.$$

A invariant set $\Sigma \subset M$ is *hyperbolic* if there exist a non-trivial measurable splitting $T_\Sigma M = E \oplus F$ and two measurable functions $0 < \mu < 1$ and $A > 0$ on Σ such that for all $x \in \Sigma$ and $n \geq 1$, we have

$$\begin{aligned} \|D\Phi^n|_{E(x)}\| &\leq A(x)\mu(x)^n \\ \|D\Phi^{-n}|_{F(\Phi^n x)}\| &\leq A(x)\mu(x)^n. \end{aligned}$$

If the functions μ and A can be replaced by constants and the splitting is continuous, then Σ is called *uniformly hyperbolic*, otherwise it is called *non-uniformly hyperbolic*. If a set is uniform hyperbolicity, then clearly it admits a dominated splitting.

3.1. Dominated splitting. In [21], Markarian, Pujals and Sambarino proved that for every contracting reflection law with $f' > 0$ and a large family of planar domains, including polygons, the invariant sets of Φ have dominated splitting. Their proof is based on the construction of an invariant cone field. We give below a proof of this fact only restricted to polygonal billiards but including reflection laws with $f' < 0$ that is much more direct than the proof in [21], not requiring the construction of an invariant cone field.

Proposition 3.1. *Every invariant set Σ of Φ has a dominated splitting. Moreover,*

- (1) *if $\limsup_{n \rightarrow +\infty} n^{-1} \log \alpha_n(x) > 0$ for every $x \in \Sigma$, then Σ is hyperbolic,*
- (2) *if there exist $A > 0$ and $\mu > 1$ such that $\alpha_n(x) \geq A\mu^n$ for every $x \in \Sigma$, then Σ is uniformly hyperbolic.*

Proof. Suppose that Σ is an invariant set. Given a point $x_0 \in \Sigma$, let $x_n = \Phi(x_0)$. From (2.2), we obtain

$$D\Phi^{-n}(x_n) = (-1)^n \begin{pmatrix} \alpha_n^{-1}(x_0) & -\frac{\gamma_n(x_0)}{\alpha_n(x_0)\beta_n(x_0)} \\ 0 & \beta_n^{-1}(x_0) \end{pmatrix} \quad \text{for } n > 0.$$

Let $F = \{F(x_0)\}_{x_0 \in \Sigma}$ be the horizontal subbundle on Σ given by $F(x_0) = \{u \in T_{x_0} M : u_\theta = 0\}$. Clearly F is continuous and invariant.

Also, we have

$$\|D\Phi^{-n}|_{F(x_n)}\| = \alpha_n^{-1}(x_0) \quad \text{for } n > 0. \quad (3.1)$$

Next, let $V(x_0) = \{u \in T_{x_0}M : u_s = 0\}$ be the vertical subspace at $x_0 \in \Sigma$. We define the subbundle E as the collection $\{E(x_0)\}_{x_0 \in \Sigma}$ with $E(x_0)$ being the subspace of $T_{x_0}M$ defined by

$$E(x_0) = \lim_{n \rightarrow +\infty} D\Phi^{-n}(x_n)V(x_n), \quad (3.2)$$

where the limit has to be understood in the projective sense. Since

$$D\Phi^{-n}(x_n) \begin{pmatrix} 0 \\ 1 \end{pmatrix} = -(\beta_n(x_0))^{-n} \begin{pmatrix} -\frac{\gamma_n(x_0)}{\alpha_n(x_0)} \\ 1 \end{pmatrix},$$

the existence of the limit in (3.2) follows from the convergence of the sequence $\{\gamma_n(x_0)/\alpha_n(x_0)\}_{n \geq 1}$. As a matter of fact, this sequence is uniformly absolutely-convergent on Σ . Indeed,

$$\lim_{n \rightarrow +\infty} \frac{\gamma_n(x_0)}{\alpha_n(x_0)} = \frac{1}{\cos \theta_0} \sum_{i=0}^{\infty} \frac{t(x_i)}{\Lambda_i(x_1)} \beta_i(x_i),$$

and since $t(x_i) \leq \text{diam } P$ and $1 \leq \Lambda_i(x_1)$ for every i , $|\theta_0| < \lambda\pi/2$ by Lemma 2.3, and $|\beta_n| \leq \lambda^n$, we have

$$\frac{1}{\cos \theta_0} \sum_{i=0}^{\infty} \frac{t(x_i)}{\Lambda_i(x_1)} |\beta_i(x_i)| \leq \frac{1}{\cos \theta_0} \sum_{i=0}^{\infty} \frac{t(x_i)}{\Lambda_i(x_1)} \lambda^i \leq \left(\cos \frac{\pi\lambda}{2} \right)^{-1} \frac{\text{diam } P}{1 - \lambda}.$$

From the convergence of $\{\gamma_n(x_0)/\alpha_n(x_0)\}_{n \geq 1}$, we can immediately deduce that the subspaces $E(x_0)$ and $F(x_0)$ are transversal, and the subbundle E is invariant (since $E(x_0)$ is defined as a limit). Moreover, from the uniform absolute convergence of $\{\gamma_n(x_0)/\alpha_n(x_0)\}_{n \geq 1}$ and the continuity of functions $\gamma_n(x_0)/\alpha_n(x_0)$ on Σ , we deduce that E is continuous on Σ .

Now, it is not difficult to see that

$$\begin{aligned} \|D\Phi^n|_{E(x_0)}\| &= \lim_{k \rightarrow +\infty} \frac{\|D\Phi^{n-k}(x_k)(0, 1)^T\|}{\|D\Phi^{-k}(x_k)(0, 1)^T\|} \\ &= \lambda^n \lim_{k \rightarrow +\infty} \frac{\left(1 + \left(\frac{\gamma_{k-n}(x_n)}{\alpha_{k-n}(x_n)}\right)^2\right)^{\frac{1}{2}}}{\left(1 + \left(\frac{\gamma_k(x_0)}{\alpha_k(x_0)}\right)^2\right)^{\frac{1}{2}}} \\ &\leq \lambda^n \left(1 + \left(\cos \frac{\pi\lambda}{2}\right)^{-1} \frac{\text{diam } P}{1 - \lambda}\right). \end{aligned}$$

It follows that $\|D\Phi^n|_{E(x_0)}\| \leq A\lambda^n$ for some constant $A > 0$ independent of x_0 and n . This means that the subbundle E is uniformly

contracting. Now, using (3.1) and the fact that $1/\alpha_n(x_0)$ is uniformly bounded by $1/\cos(\pi\lambda/2)$, we can conclude that

$$\|D\Phi^n|_{E(x_0)}\| \cdot \|D\Phi^{-n}|_{F(x_n)}\| \leq \frac{A}{\cos(\pi\lambda/2)} \lambda^n \quad \text{for } x_0 \in \Sigma \text{ and } n \geq 1,$$

i.e., the invariant set Σ has dominated splitting.

Finally, to prove (1) and (2), we just need to observe that the subbundle E is uniformly contracting, and that under the hypotheses of (1) and (2), the subbundle F is expanding and uniformly expanding, respectively. \square

3.2. Periodic orbits.

Definition 3.1. We say that $K \subset M$ is an *attracting set* if K has a neighborhood U in M such that $d(\Phi^n(x), K) \rightarrow 0$ as $n \rightarrow +\infty$ for every $x \in U \setminus N_\infty^+$.

Definition 3.2. We denote by \mathcal{P} the set of all periodic points of period two of the billiard map Φ .

Proposition 3.2. *The periodic points of Φ have the following properties:*

- (1) *every periodic point of period two is parabolic,*
- (2) *the set \mathcal{P} is an attracting set,*
- (3) *every periodic point of period greater than two is hyperbolic.*

Proof. Suppose that (s_0, θ_0) is a periodic point of Φ of period n . Since $(s_n, \theta_n) = (s_0, \theta_0)$, it follows immediately from (2.3) that $\alpha_n(s_0, \theta_0) = \Lambda_n(s_0, \theta_0)$. If we define $\zeta = \max\{\rho(\bar{\theta}_i) : i = 1, \dots, n\}$, then we see that $\zeta \leq \alpha_n(s_0, \theta_0) \leq \zeta^n$.

Since a periodic point (s_0, θ_0) of period two corresponds to a billiard trajectory that hits perpendicularly two sides of P , we have $\theta_0 = \theta_1 = \theta_2 = 0$. Hence $\zeta = 1$, and so $\alpha_2(s_0, \theta_0) = 1$. As $(-1)^n \alpha_n$ is an eigenvalue of $D\Phi^n$ (cf. (2.2)), all periodic points of period two are parabolic. Using the fact that f is a contraction, it is not difficult to show that the set of all periodic points of period two is an attracting set.

Finally, we show that periodic points of period $n > 2$ are hyperbolic. These points have at least two collisions with a non-zero angle, and so $\rho(\bar{\theta}_i) > 1$ for at least two i 's. If we set $\mu = \zeta^{1/n} > 1$, then $\alpha_n(s_0, \theta_0) > \mu^n$. The wanted result now follows from part (2) of Proposition 3.1. \square

3.3. Uniform hyperbolicity. We now address the problem of the hyperbolicity of general invariant sets. Proposition 3.3 below represents a considerable improvement of a result of Markarian, Pujals and Sambarino, stating that for any convex polygon without parallel sides, the map Φ_f is hyperbolic provided that $\lambda(f)$ is sufficiently close to zero [21, Corollary 4 and Theorem 23].

Definition 3.3. We say that a forward invariant set $\Omega \subset M$ has Property (A) if there exists $m > 0$ such that every sequence of consecutive collisions between parallel sides of P contained in Ω consists of no more than m collisions.

Proposition 3.3. *Suppose that Σ is an invariant set with Property (A). Then Σ is uniformly hyperbolic.*

Proof. Let $\Delta = \min_{i,j} |\pi - \delta(L_i, L_j)|/2$, where the minimum is taken over all pairs of non-parallel sides L_i and L_j seeing each other. It is easy to check that $\Delta > 0$.

Let $\{(s_n, \theta_n)\}_{n \in \mathbb{Z}}$ be an orbit contained in Σ . Consider two consecutive collisions (s_i, θ_i) and (s_{i+1}, θ_{i+1}) at non-parallel sides of P . By (2.1), we have

$$\bar{\theta}_{i+1} = \pi - \delta - \theta_i,$$

where δ is the angle formed by the sides of P containing s_i and s_{i+1} . Fix $0 < \epsilon < \Delta$. If $|\theta_i| \geq \epsilon$, then $|\bar{\theta}_i| > \epsilon$ so that $\rho(\bar{\theta}_i) \geq r(\epsilon)$. On the other hand, if $|\theta_i| < \epsilon$, then $|\bar{\theta}_{i+1}| > \pi - \delta - \epsilon > 2\Delta - \epsilon > \epsilon$, and so $\rho(\bar{\theta}_{i+1}) \geq r(\epsilon)$. Since $\rho \geq 1$, in both cases, we have

$$\rho(\bar{\theta}_i)\rho(\bar{\theta}_{i+1}) \geq r(\epsilon). \quad (3.3)$$

By Property (A), the maximum number of consecutive collisions occurring at parallel sides of P must be bounded above by $m > 0$. If P does not have pair of parallel sides, then we set $m = 1$. Now, consider a sequence of $m+2$ consecutive collisions $(s_j, \theta_j), \dots, (s_{j+m+1}, \theta_{j+m+1})$. It is clear that such a sequence contain at least one pair of consecutive collisions occurring at sides that are not parallel. Hence, by using (3.3) and $\rho \geq 1$, we deduce that

$$\prod_{k=0}^{m+1} \rho(\bar{\theta}_{j+k}) \geq r(\epsilon).$$

From this inequality, we easily obtain $\Lambda_n(s_1, \theta_1) \geq r(\epsilon)^{\frac{n-m-1}{m+2}}$ for all $n \geq m+2$. But $\Lambda_n \geq 1$ for every $n \geq 1$ and so

$$\Lambda_n(s_1, \theta_1) \geq r(\epsilon)^{\frac{n-m-1}{m+2}} \quad \text{for } n \geq 1. \quad (3.4)$$

Define $A = r(\epsilon)^{-1} \cos(\pi\lambda/2)$ and $\mu = r(\epsilon)^{1/(m+2)}$. Combining (2.3) and (3.4), and using Lemma 2.3, we obtain

$$\alpha_n(s_0, \theta_0) \geq A\mu^n \quad \text{for } n \geq 1.$$

Therefore, Φ is uniformly expanding along the horizontal direction. By Part (2) of Proposition 3.1, it follows that Σ is uniformly hyperbolic. \square

Corollary 3.4. *Suppose that $\mathcal{P} = \emptyset$. Then D is uniformly hyperbolic.*

Proof. Since $\mathcal{P} = \emptyset$, we can apply Proposition 3.3 to D . \square

All regular $(2n+1)$ -gons satisfy the hypothesis of Corollary 3.4.




FIGURE 3. A uniformly hyperbolic polygonal billiard with parallel sides.

Corollary 3.5. *The set D is uniformly hyperbolic for every regular $(2n + 1)$ -gon.*

Remark 3.6. A necessary condition for $\mathcal{P} \neq \emptyset$ is that P has a pair of parallel sides. This condition is clearly not sufficient. Fig. 3 displays an example of a polygon with parallel sides satisfying the hypothesis of Corollary 3.4.

Even when $\mathcal{P} \neq \emptyset$, the map Φ may still have uniformly hyperbolic sets. In Section 8, we will show that this is the case for billiards in rectangles with proper contracting reflection laws (see Proposition 8.8).

4. SRB MEASURES FOR HYPERBOLIC ATTRACTORS

In this section we find criteria for the existence of SRB measures in polygonal billiards with a contracting reflection law. This is a consequence of a general result of Pesin that states the existence of such measures for piecewise smooth systems satisfying certain technical conditions. We show that those holds for the case of the billiards consider.

Recall that an invariant probability measure μ of a diffeomorphism $g: U \rightarrow U$ on a Riemannian manifold U is called an *SRB measure* (after Sinai, Ruelle and Bowen) if it is hyperbolic, and its conditional measures on unstable manifolds are absolutely continuous. The relevance of such a measure is due to the fact that if μ is also ergodic, then there exists a positive Lebesgue measure set of points $x \in U$ such that

$$\lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{i=0}^{n-1} \varphi(g^i(x)) = \int_U \varphi d\mu$$

for every continuous function $\varphi: U \rightarrow \mathbb{R}$. For the setting considered in this section, this property is proved in [23, Theorem 3]. We refer the reader to [36], for a nice review on SRB measures.

4.1. Piecewise smooth maps. Let U be a smooth Riemannian surface with metric ρ and area ν . Let K be an open connected subset of U with compact closure.

Definition 4.1. We say that g is a *piecewise smooth map* on K if there exist four constants $a_1, a_2 \geq 0$, $c_1, c_2 > 0$ and finitely many disjoint connected open sets $\{K_i\}$ with boundary formed by a finite number of C^1 curves such that $\overline{K} = \bigcup_i \overline{K_i}$, $g: \bigcup_i K_i \rightarrow U$ is injective, $g|_{K_i}$ is a C^2 diffeomorphism on its image satisfying

$$\begin{aligned} \|D^2 g(x)\| &\leq c_1 \rho(x, N)^{-a_1}, & x \in K \setminus N, \\ \|D^2 g^{-1}(x)\| &\leq c_2 \rho(x, N')^{-a_2}, & x \in K \setminus N', \end{aligned} \quad (4.1)$$

where $N = \bigcup_i \partial K_i$ and $N' = \partial(g(K \setminus N))$.

Next, we define

$$K^+ = \{x \in K : g^n(x) \notin N, n \geq 0\},$$

and consider the invariant set

$$D = \bigcap_{n \geq 0} g^n(K^+).$$

We call the closure $A = \overline{D}$ an *attractor* following the nomenclature of [23]. If D is uniformly hyperbolic, then we refer to A as a generalized hyperbolic attractor, or simply *hyperbolic attractor*. The set A is said to be *regular* if

$$D_l^- = \bigcup_{0 < \delta \leq 1} \{x \in A : \rho(g^{-n}(x), N') \geq \delta e^{-nl}, n \geq 0\}$$

is not empty for all sufficiently small $l > 0$. In particular, A is regular if it contains a periodic orbit. Moreover, [23, Proposition 3] states that if there are $c > 0$ and $q > 0$ such that for every $n \geq 0$ and every $\varepsilon > 0$,

$$\nu(K^+ \cap g^{-n}(N_\varepsilon)) \leq c \varepsilon^q, \quad (4.2)$$

then A is regular. Here N_ε denotes the ε -neighborhood of N .

If A is a regular hyperbolic attractor, then there is $l > 0$ such that any $x \in D^- := D_l^-$ has local unstable manifold $W_{loc}^u(x)$ [23, Proposition 4]. The next theorem provides sufficient conditions for the existence of SRB measures supported on A .

Theorem 4.1 ([23]). *Suppose that*

- (1) *A is a regular hyperbolic attractor,*
- (2) *there exist $x \in D^-$ and $c, q, \varepsilon_0 > 0$ such that for $n \geq 0$ and $0 < \varepsilon < \varepsilon_0$, we have*

$$\ell(W_{loc}^u(x) \cap g^{-n}(N_\varepsilon)) \leq c \varepsilon^q, \quad (4.3)$$

where ℓ is the length of a curve in U .

Then A has countably many ergodic SRB measures supported on D .

Under more restrictive conditions, which for polygonal billiard maps follow from Proposition 4.3 below, Sataev proved that the number of ergodic SRB measures is actually finite [25]. These measures have strong ergodic properties: for every ergodic SRB measure μ , there exist $n > 0$

and a g^n -invariant subset E with $\mu(E) > 0$ such that $(g^n|_E, \mu|_E)$ is Bernoulli and has exponential decay of correlations, and the Central Limit Theorem holds for $(g^n|_E, \mu|_E)$ [2, 10, 35]. Moreover, the Young dimension formula, relating the Hausdorff dimension of μ to its Lyapunov exponents and entropy, holds true [27]. Notice also that under the conditions of the above theorem, the hyperbolic points are dense in the attractor [23, Theorem 11].

Known examples of the above systems are Lozi maps, Belykh maps and geometrical Lorenz maps [1, 6, 20, 34]. We will show in the following that polygonal billiards with strongly contracting reflection laws are also of this type.

4.2. SRB measures for polygonal billiards. Let $K = M$ for the polygonal billiard map Φ_f where f is any reflection law, and $N = N_1^+$. We recall that $N_1^+ = \partial M \cup V \cup S_1^+$. From Proposition 2.1, it follows that the set N_1^+ divides M into pairwise disjoint open connected subsets M_1, \dots, M_k , i.e., $M \setminus N_1^+ = \bigcup_i M_i$. We denote by $\Phi_{f,i}$ the restriction $\Phi_f|_{M_i}$.

The map Φ_f for a contracting reflection law $f \in \mathcal{R}_1^k$, $k \geq 1$, is a C^k -diffeomorphism from $M \setminus N_1^+$ onto its image in M .

The set of points that can be infinitely iterated is $K^+ = M \setminus N_\infty^+$ and let

$$A = \overline{\bigcap_{n \geq 0} \Phi_f^n(K^+)}.$$

Notice that A contains for example the period two orbits, if they exist.

We say that a polygon has *parallel sides facing each other* if there is a straight line joining orthogonally two of its sides. This is the same as the billiard map having a period two orbit (i.e. $\mathcal{P} \neq \emptyset$).

Proposition 4.2. *A is a hyperbolic attractor iff the polygon does not have parallel sides facing each other.*

Proof. A is a hyperbolic attractor iff it does not contain the parabolic orbits that are the period two orbits (Proposition 3.3). These exist iff there are parallel sides facing each other. \square

In the following we will prove the conditions of Theorem 4.1 for the billiard map Φ_f .

A horizontal curve Γ is a curve lying on a line of constant θ inside M_i for some i . It is parametrized by a smooth path

$$\gamma: I \rightarrow M_i \tag{4.4}$$

on an interval I contained in $[0, 1]$, satisfying $\gamma'(t) = (1, 0)$, $t \in I$. So, $\ell(\Gamma) = |I| \leq 1$.

Given $n \geq 1$, let $p(S_n^+)$ be the maximum number of smooth components of S_n^+ intersecting at one point. Moreover, denote the smallest

expansion rate of Φ_f^n along the unstable direction by

$$\alpha(\Phi_f^n) = \inf_{x \in M \setminus N_n^+} \|D\Phi_f^n(x)(1, 0)\|.$$

Notice that $\alpha(\Phi_f^n) \geq \alpha(\Phi_f)^n$ and that $f \mapsto \alpha(\Phi_f)$ is a continuous map because N_1^+ does not depend on f .

To apply Theorem 4.1 to polygonal billiards one needs to prove (4.2) and (4.3). Both conditions follow from the conclusion (4.6) of Proposition 4.3 below. This proposition is the adaptation of [23, Theorem 14] (see also [27, Theorem 6.1] and [35, Section 7]) to polygonal billiards. This adaptation is necessary because billiards do not satisfy all conditions of the mentioned results (e.g. existence of smooth extensions) and verify trivially others (like bounded distortion estimates).

Proposition 4.3. *If $m \geq 1$ and*

$$p(S_m^+) < \alpha(\Phi_f^m), \quad (4.5)$$

then there exist $c_m, \varepsilon_0 > 0$ such that for any horizontal curve $\Gamma \subset \tilde{D}$, $n \geq 0$ and $0 < \varepsilon < \varepsilon_0$,

$$\ell(\Gamma \cap \Phi_f^{-n}(N_\varepsilon)) \leq c_m \varepsilon. \quad (4.6)$$

Proof. Notice that by considering $\varepsilon_0 < (1 - \lambda)\pi/2$, the orbit of any point never enters the ε_0 -neighborhood of ∂M . So N_ε in Condition (4.6) can be safely replaced by the ε -neighborhood of $V \cup S_1^+$. Accordingly, throughout this proof, the symbols N_1^+ and N_ε have to be understood as $V \cup S_1^+$ and the ε -neighborhood of $V \cup S_1^+$, respectively.

We have that $G_r = \Gamma \cap N_r^+$ is finite for any $r \geq 0$, where we set $N_0^+ = \emptyset$. Indeed, after one iteration of the billiard map only a finite number of points in the horizontal curve Γ reach a corner, and the same after r iterations for each of the components of the image of Γ . So, $\ell(\Gamma) = \ell(\Gamma \setminus G_r) \leq 1$ and

$$\begin{aligned} \ell(\Gamma \cap \Phi_f^{-r}(N_\varepsilon)) &= \ell((\Gamma \setminus G_r) \cap \Phi_f^{-r}(N_\varepsilon)) \\ &= \ell(\Phi_f^{-r}(\Phi_f^r(\Gamma \setminus G_r) \cap N_\varepsilon)). \end{aligned}$$

The r -th iterate of $\Gamma \setminus G_r$ is the union of a finite number of smooth curves (each one diffeomorphic to a component of $\Gamma \setminus G_r$) given by

$$\Phi_f^r(\Gamma \setminus G_r) = \bigcup_{i \in J_r} \Gamma_{r,i},$$

where J_r is finite. We have that $\Gamma_{r,i} = \Phi_f(\Gamma_{r-1,j} \cap M_l)$ for some j, l depending on i, r and $j \in J_{r-1}$. Notice that the set J_r does not have more elements than J_{r+1} , and that $\Gamma_{0,i} = \Gamma$ where $i \in J_0 = \{1\}$.

Fix $n = sm + s'$ where $s \geq 0$ and $s' \in \{0, \dots, m-1\}$. Since all $\tilde{\Gamma}_{n,i} = \Gamma_{n,i} \cap N_\varepsilon$ are disjoint,

$$\ell(\Gamma \cap \Phi_f^{-n}(N_\varepsilon)) = \ell\left(\Phi_f^{-sm} \circ \Phi_f^{-s'}\left(\bigcup_{i \in J_n} \tilde{\Gamma}_{n,i}\right)\right) = \sum_{i \in J_n} \ell\left(\Phi_f^{-sm}(\tilde{\Gamma}'_{sm,i})\right)$$

where $\tilde{\Gamma}'_{sm,i} = \Phi_f^{-s'}(\tilde{\Gamma}_{n,i})$.

From the transversality between N_1^+ and the horizontal direction given by Proposition 2.1, there is $C' > 0$ (independent of n and Γ) satisfying the inequality $\ell(\tilde{\Gamma}_{n,i}) \leq C'\varepsilon$. Let η be the maximum number of intersections between a horizontal line in the phase space M and N_1^+ . Given $j \in J_{sm}$ denote by $J_{n,j}$ the set of all $i \in J_n$ satisfying $\tilde{\Gamma}'_{sm,i} \subset \Gamma_{sm,j}$. Thus, $\#J_{n,j} \leq (\eta + 1)^{s'}$. Notice that $\cup_{j \in J_{sm}} J_{n,j} = J_n$. Furthermore,

$$\sum_{i \in J_{n,j}} \ell(\tilde{\Gamma}'_{sm,i}) \leq \sum_{i \in J_{n,j}} \frac{\ell(\tilde{\Gamma}_{n,i})}{\alpha(\Phi_f^{s'})} \leq \frac{(\eta + 1)^m C' \varepsilon}{\min_{s'' < m} \alpha(\Phi_f^{s''})} = C'_m \varepsilon. \quad (4.7)$$

Notice that the definition of $p = p(S_m^+)$ is equivalent to saying that it is the smallest positive integer for which there is $C_m > 0$ such that if Γ is a horizontal curve with $\ell(\Gamma) < C_m$, then $\Gamma \setminus S_m^+$ has at most p components with positive length.

Let $d = \min\{C_m, \ell(\Gamma)\}$. Write $\psi = \Phi_f^m$ and $\Upsilon_{r,i} = \Gamma_{rm,i}$ with $i \in J_r$ and $0 \leq r \leq s$.

Take any $0 \leq k \leq s$ and $l \in J_{km}$ such that $\ell(\Upsilon_{k,l}) \geq d$, denote the set of these pairs (k, l) by \mathcal{I}_s and the set of indices

$$I(k, l) = \{i \in J_{sm} : \psi^{-(s-k)}(\Upsilon_{s,i}) \subset \Upsilon_{k,l}\}.$$

Now, define

$$I_{k,l} = I(k, l) \setminus \bigcup_{(r,j) \in \mathcal{I}_s, r > k} I(r, j).$$

It is now easy to check that the sets $I_{k,l}$ are disjoint, their union is J_{sm} and $\#I_{k,l} \leq p^{s-k-1}$. Therefore,

$$J_n = \bigcup_{(k,l) \in \mathcal{I}_s} \bigcup_{j \in I_{k,l}} J_{n,j}.$$

For each $(k, l) \in \mathcal{I}_s$ define

$$\Lambda_{k,l} = \bigcup_{j \in I_{k,l}} \bigcup_{i \in J_{n,j}} \psi^{-(s-k)}(\tilde{\Gamma}'_{sm,i}) \subset \Upsilon_{k,l}.$$

Notice that this is a disjoint union, hence

$$\begin{aligned} \sum_{i \in J_n} \ell\left(\Phi_f^{-sm}(\tilde{\Gamma}'_{sm,i})\right) &\leq \sum_{(k,l) \in \mathcal{I}_s} \sum_{j \in I_{k,l}} \sum_{i \in J_{n,j}} \ell\left(\psi^{-s}(\tilde{\Gamma}'_{sm,i})\right) \\ &= \sum_{(k,l) \in \mathcal{I}_s} \ell\left(\psi^{-k}(\Lambda_{k,l})\right). \end{aligned}$$

Each $\Upsilon_{k,l}$ is parametrized by the path $\gamma_{k,l}(t) = \psi^k \circ \gamma(t)$ with $\gamma'_{k,l}(t) = D\psi^k \circ \gamma(t)(1, 0)$ and

$$\ell(\Upsilon_{k,l}) = \int_{\gamma_{k,l}^{-1}(\Upsilon_{k,l})} \|\gamma'_{k,l}(t)\| dt = \|\gamma'_{k,l}(\xi)\| \ell(\psi^{-k}(\Upsilon_{k,l})),$$

where $\xi \in \gamma_{k,l}^{-1}(\Upsilon_{k,l}) \subset [0, 1]$. Similarly, $\ell(\Lambda_{k,l}) = \|\gamma'_{k,l}(\xi')\| \ell(\psi^{-k}(\Lambda_{k,l}))$, for some $\xi' \in \gamma_{k,l}^{-1}(\Lambda_{k,l})$. So,

$$\ell(\psi^{-k}(\Lambda_{k,l})) = \ell(\Lambda_{k,l}) \frac{\|\gamma'_{k,l}(\xi)\|}{\|\gamma'_{k,l}(\xi')\|} \frac{\ell(\psi^{-k}(\Upsilon_{k,l}))}{\ell(\Upsilon_{k,l})}. \quad (4.8)$$

Following Section 2.5, we have that $\|D\psi^k(x)(1, 0)\|$ is constant in $\psi^{-k}(\Upsilon_{k,l}) \subset \Gamma$. This proves that $\|\gamma'_{k,l}(\xi)\| = \|\gamma'_{k,l}(\xi')\|$. Moreover, from (4.7), writing $\alpha = \alpha(\psi)$ which is larger than p by (4.5),

$$\ell(\Lambda_{k,l}) \leq \sum_{j \in I_{k,l}} \sum_{i \in J_{n,j}} \frac{\ell(\tilde{\Gamma}'_{sm,i})}{\alpha^{s-k}} \leq C'_m \varepsilon \left(\frac{p}{\alpha}\right)^{s-k}.$$

Finally, from $\ell(\Upsilon_{k,l}) \geq d$ when $(k, l) \in \mathcal{I}_s$, using (4.8),

$$\sum_{(k,l) \in \mathcal{I}_s} \ell(\psi^{-k}(\Lambda_{k,l})) \leq \frac{C'_m \varepsilon}{d} \sum_{k=0}^s \left(\frac{p}{\alpha}\right)^{s-k} \sum_l \ell(\psi^{-k}(\Upsilon_{k,l})).$$

Furthermore, $\sum_l \ell(\psi^{-k}(\Upsilon_{k,l})) \leq \ell(\psi^{-k}(\cup_l \Upsilon_{k,l})) = \ell(\Gamma)$.

All the above estimates imply that

$$\ell(\Gamma \cap \Phi_f^{-n}(N_\varepsilon)) \leq \frac{C'_m \alpha \ell(\Gamma)}{d(\alpha - p)} \varepsilon.$$

The fact that $d^{-1}\ell(\Gamma)$ is equal to 1 if $\ell(\Gamma) \leq C_m$ and otherwise bounded from above by C_m^{-1} because $\ell(\Gamma) \leq 1$, proves (4.6). \square

Denote by \mathcal{B} the subset of contractive C^2 reflection laws f in \mathcal{R}_1^2 such that f and its inverse f^{-1} have bounded second derivatives.

Theorem 4.4. *Consider a polygon without parallel sides facing each other and suppose that $f \in \mathcal{B}$. If there is $m \geq 1$ such that*

$$p(S_m^+) < \alpha(\Phi_f^m),$$

then A has countably many ergodic SRB measures.

Proof. First, notice that by Proposition 4.2 A is a generalized hyperbolic attractor. Moreover, the billiard map Φ_f satisfies conditions (4.1). This is a consequence of the following facts: $\Phi_f = R_f \circ \bar{\Phi}$, the conservative billiard map $\bar{\Phi}$ verifies conditions (4.1) with $a_1 = a_2 = 3$ (see [18]) and R_f has bounded second derivatives.

Now, the existence of countably many ergodic SRB measures follows from Theorem 4.1 provided that (4.2) and (4.3) hold. Both conditions follow from (4.6); the second one directly, whereas the first one after decomposing K^+ into horizontal curves. \square

Remark 4.5. In fact, (4.6) allows us to use a result by Sataev [25, Theorem 5.15], and conclude that the number of ergodic SRB measures is finite.

Remark 4.6. The conditional measures of the SRB measures in Theorem 4.4 coincide up to a normalizing constant factor with the measure induced by the Riemannian metric on the unstable manifolds. This can be deduced easily from the computations in the proof of [23, Proposition 7]. In particular, note that the function κ in that proof is identically equal to one for our billiards.

4.3. Sufficient conditions. A convex polygonal billiard corresponds to $p(S_1^+) = 2$, because in this case S_1^+ is a disjoint union of smooth curves. A non-convex polygonal billiard yields instead $\underline{p}(S_1^+) \geq 2$.

Every $\Phi_{f,i}$ admits a continuous extension map to \overline{M}_i denoted by $\hat{\Phi}_{f,i}$, which can be multi-valued at a finite number of points (these correspond to trajectories of the billiard flow tangent to the sides of the polygon). This allows us to extend Φ_f to N_1^+ in the following multi-valued way,

$$\Phi_f^n(B) = \bigcup_i \hat{\Phi}_{f,i}(\Phi_f^{n-1}(B) \cap \overline{M}_i), \quad B \subset M, \quad n \geq 1, \quad (4.9)$$

and $\Phi_f^0 = \text{id}$ on M . Similarly, for the pre-images,

$$\Phi_f^{-n}(B) = \bigcup_i \hat{\Phi}_{f,i}^{-1}(\Phi_f^{-(n-1)}(B)). \quad (4.10)$$

Lemma 4.7.

- (1) For any $n \geq 1$ and $B \subset M$, $\Phi_f^n(B) \cap B = \emptyset$ is equivalent to $\Phi_f^{-n}(B) \cap B = \emptyset$.
- (2) If there is $n \geq 1$ and $B \subset M$ such that $\Phi_f^k(B) \cap B = \emptyset$ for any $1 \leq k \leq n$, then

$$\Phi_f^{-r}(B) \cap \Phi_f^{-s}(B) = \emptyset,$$

where $1 \leq r - s \leq n$.

Proof. From $\Phi_f^n(B) \cap B = \emptyset$ we have that, for any i_1 ,

$$\hat{\Phi}_{f,i_1}(\Phi_f^{n-1}(B) \cap \overline{M}_{i_1}) \cap B = \emptyset.$$

Then its pre-image by $\hat{\Phi}_{f,i_1}$ is still empty, i.e.

$$\Phi_f^{n-1}(B) \cap \overline{M}_{i_1} \cap \hat{\Phi}_{f,i_1}^{-1}(B) = \emptyset. \quad (4.11)$$

The reverse is also true since (4.11) implies that the image of $\Phi_f^{n-1}(B) \cap \overline{M}_{i_1}$ by $\hat{\Phi}_{f,i_1}$ can not intersect B . Using the fact that $\hat{\Phi}_{f,i_1}^{-1}(B) \subset \overline{M}_{i_1}$, we have shown that $\Phi_f^n(B) \cap B = \emptyset$ is equivalent to $\Phi_f^{n-1}(B) \cap \hat{\Phi}_{f,i_1}^{-1}(B) = \emptyset$ for every i_1 .

Repeating this procedure $n - 1$ more times, we get for any i_1, \dots, i_n ,

$$B \cap \hat{\Phi}_{f,i_n}^{-1} \dots \hat{\Phi}_{f,i_1}^{-1}(B) = \emptyset.$$

Finally, the union of the above sets is again empty and, by noticing that

$$\Phi_f^{-n}(B) = \bigcup_{i_1, \dots, i_n} \hat{\Phi}_{f,i_n}^{-1} \dots \hat{\Phi}_{f,i_1}^{-1}(B),$$

the proof of the first claim is complete.

By the above, $\Phi_f^{-k}(B) \cap B = \emptyset$. Applying Φ_f^{-s} we obtain that $\Phi_f^{-(k+s)}(B) \cap \Phi_f^{-s}(B) = \emptyset$. Write now $r - s = k$ to get the second claim. \square

Below we use the notation $S_n^+(f)$ to highlight the dependence on f for $n \geq 2$. The hypothesis of the next proposition means that there are no trajectories connecting two vertexes (not necessarily distinct) of the polygon formed by $(n - 1)$ straight segments.

Proposition 4.8. *Let $f_0 \in \mathcal{R}_1^1$ or $f_0 = 0$, and $n \geq 1$. If*

$$\Phi_{f_0}^k(S_1^+) \cap S_1^+ = \emptyset \quad \text{for } 1 \leq k \leq n - 1,$$

then there is $\delta > 0$ such that for $f \in \mathcal{R}_1^1$ satisfying $\lambda(f - f_0) < \delta$ we have $p(S_n^+(f)) = p(S_1^+)$.

Proof. Let $f_0 \in \mathcal{R}_1^1$. By Lemma 4.7 all pre-images of S_1^+ under Φ_{f_0} up to the iterate $n - 1$ are disjoint. Moreover, each $\Phi_{f_0,i}^{-1}(S_1^+) \subset M_i$ is diffeomorphic to connected components of S_1^+ and the same holds up to the $(n - 1)$ -th iterate. Since $f \mapsto \Phi_{f,i}$ and $f \mapsto \Phi_{f,i}^{-1}$ are continuous maps for each i , the same properties hold for f with $\lambda(f - f_0)$ small enough. Recall also the fact that each $\Phi_{f,i}^{-1}$ preserves horizontal lines. Therefore, the number $p(S_k^+(f))$ is the same for every $1 \leq k \leq n$.

Suppose now that $f_0 = 0$. In this case each $\Phi_{0,i}$ is not a diffeomorphism. For the same reasons as above, given any $f \in \mathcal{R}_1^1$ such that $\lambda(f)$ is small, $\Phi_f^k(S_1^+) \cap S_1^+ = \emptyset$, $1 \leq k \leq n - 1$. The proof follows from the previous case. \square

Theorem 4.9. *Consider a polygon without parallel sides facing each other and $f_0 \in \mathcal{R}_1^1$ or $f_0 = 0$ such that*

$$\Phi_{f_0}^k(S_1^+) \cap S_1^+ = \emptyset \quad \text{for } k \geq 1.$$

Then there is $\delta > 0$ such that for $f \in \mathcal{B}$ satisfying $\lambda(f - f_0) < \delta$, the billiard map Φ_f has a hyperbolic attractor with countably many ergodic SRB measures.

Proof. By the uniform hyperbolicity of Φ_f (see Proposition 3.3), there is some $m \geq 1$ such that $\alpha(\Phi_f^m) > 1$. Take the supremum of such m in some neighborhood of f_0 . Let $n \geq 1$ such that $\alpha(\Phi_f^{nm}) \geq \alpha(\Phi_f^m)^n > p(S_1^+)$. By Proposition 4.8 we have that $p(S_{nm}^+) = p(S_1^+)$ by choosing a smaller δ . The claim follows from Theorem 4.4. \square

Theorem 4.10. *Consider a polygon without parallel sides facing each other, $f_0 \in \mathcal{R}_1^1$ or $f_0 = 0$, and*

$$n > \frac{\log p(S_1^+)}{\log \alpha(\Phi_{f_0})}$$

such that

$$\Phi_{f_0}^k(S_1^+) \cap S_1^+ = \emptyset \quad \text{for } 1 \leq k \leq n-1.$$

Then there is $\delta > 0$ such that for $f \in \mathcal{B}$ satisfying $\lambda(f - f_0) < \delta$, the billiard map Φ_f has a hyperbolic attractor with countably many ergodic SRB measures.

Proof. Proposition 4.8 holds for Φ_f , i.e. $p(S_n^+(f)) = p(S_1^+)$, for $\lambda(f - f_0)$ small enough. From the continuity of $f \mapsto \alpha(\Phi_f)$, for any $\varepsilon > 0$ and sufficiently small $\lambda(f - f_0)$, we have $\alpha(\Phi_f^n) \geq \alpha(\Phi_f)^n > (\alpha(\Phi_{f_0}) - \varepsilon)^n$. By the hypothesis on n we get $\alpha(\Phi_{f_0}) > p(S_1^+)^{1/n}$. So, for ε small enough (depending on n) we have $\alpha(\Phi_f^n) > p(S_1^+)$. The result now follows from Theorem 4.4. \square

The study of the ergodic properties of the SRB measures obtained in Theorems 4.9 and 4.10 will appear elsewhere [16].

5. GENERIC POLYGONS

We now introduce the moduli spaces of polygons and prove that, generically, polygons in these spaces have no “orthogonal vertex connections”. This will be used later to prove the existence of SRB measures for small $\lambda(f)$.

5.1. Moduli spaces of polygons. Any sequence $P = (v_0, v_1, \dots, v_{n-1})$ of points in \mathbb{R}^2 determines a *closed polygonal line*. The line segments $e_1 = [v_0, v_1]$, $e_2 = [v_1, v_2]$, \dots , $e_0 = e_n = [v_{n-1}, v_0]$, are called the *edges* of P . We say that P is a *n-gon* if the lines supporting e_{i-1} and e_i are always distinct, and the polygonal line P is non self-intersecting. Two *n-gons* $P = (v_0, v_1, \dots, v_{n-1})$, $P' = (v'_0, v'_1, \dots, v'_{n-1})$ are said to be *similar* if there is an orientation preserving similarity T of the Euclidean plane such that $T(v_i) = v'_i$, for $i = 0, 1, \dots, n-1$. Similarity is an equivalence relation on the space of all *n-gons*, and the quotient by this relation is called the *moduli space* of *n-gons*, denoted hereafter by \mathcal{P}_n . Let \mathbb{P}^1 and \mathbb{P}^2 denote the real projective line and plane, respectively.

Proposition 5.1. *The moduli space \mathcal{P}_n is diffeomorphic to an open semialgebraic subset of $\mathbb{P}^1 \times (\mathbb{P}^2)^{n-3} \times \mathbb{P}^1$, and it is a manifold of dimension $2n - 4$.*

Proof. Each edge e_i determines a line in the Euclidean plane \mathbb{R}^2 , defined by an affine equation $a_i x + b_i y + c_i = 0$, and we shall refer to $(a_i : b_i : c_i)$

as the projective coordinates of e_i . The vertices v_0, v_1, \dots, v_{n-1} of the polygon, $v_i = e_{i-1} \cap e_i$, are easily computed to have coordinates

$$(x_i, y_i) = \left(\frac{b_{i-1}c_i - b_i c_{i-1}}{a_{i-1}b_i - a_i b_{i-1}}, \frac{a_i c_{i-1} - a_{i-1} c_i}{a_{i-1}b_i - a_i b_{i-1}} \right). \quad (5.1)$$

We are adopting the convention that $e_0 = e_n$. Up to a similarity transformation, we can assume that $v_0 = (-1, 0)$ and $v_1 = (1, 0)$, and specify coordinates for the edges e_0, e_1 and e_{n-1} respectively of the form $(a_0 : b_0 : c_0) = (0, 1, 0)$, $(a_1 : b_1 : c_1) \equiv (a_1 : b_1)$ and $(a_{n-1}, b_{n-1}, -a_{n-1}) \equiv (a_{n-1} : b_{n-1})$. Hence any n -gon P determines and is determined by the vector in $\mathbb{P}^1 \times (\mathbb{P}^2)^{n-3} \times \mathbb{P}^1$,

$$\theta(P) = [(a_1 : b_1), (a_2 : b_2 : c_2), \dots, (a_{n-2} : b_{n-2} : c_{n-2}), (a_{n-1} : b_{n-1})].$$

The values of the projective coordinates of the edges missing in the vector $\theta(P)$ are fixed to be $(a_0 : b_0 : c_0) = (0 : 1 : 0)$, $c_1 = a_1$ and $c_{n-1} = -a_{n-1}$. Since consecutive edges are not parallel, we must have

$$a_{i-1}b_i - a_i b_{i-1} \neq 0 \quad \text{for } i = 0, 1, \dots, n-1. \quad (5.2)$$

Under these conditions, the coordinates (x_i, y_i) of the vertices in (5.1) are well defined. The second condition in the definition of a n -gon is equivalent to saying that for $i \neq j$ and $i \neq j \pm 1$ either the lines $(a_i : b_i : c_i)$ and $(a_j : b_j : c_j)$ are parallel, or else their intersection point

$$(u_{ij}, v_{ij}) = \left(\frac{b_j c_i - b_i c_j}{a_j b_i - a_i b_j}, \frac{a_i c_j - a_j c_i}{a_j b_i - a_i b_j} \right)$$

does not lie in $e_i \cup e_j$, a condition which is easily seen to be semialgebraic. Hence the subset $\Omega_n \subseteq \mathbb{P}^1 \times (\mathbb{P}^2)^{n-3} \times \mathbb{P}^1$ of all data satisfying both (5.2) and the previous condition is an open semialgebraic subset of $\mathbb{P}^1 \times (\mathbb{P}^2)^{n-3} \times \mathbb{P}^1$. The previous considerations show that the map θ induces a diffeomorphism $\theta: \mathcal{P}_n \rightarrow \Omega_n$. \square

Given a n -gon P with edges e_1, \dots, e_n , denote by ℓ_i the line supporting e_i , and by $\pi_i : \mathbb{R}^2 \rightarrow \ell_i$ the orthogonal projection onto ℓ_i . We say that P has an *orthogonal vertex connection of order m* if there is a sequence of edges $e_{i_1}, \dots, e_{i_{m'}}$, with length $m' \leq m$, and a sequence $p_0, \dots, p_{m'}$ of points in P such that:

- (a) p_0 and $p_{m'}$ are vertices of P ;
- (b) $p_k = \pi_{i_k}(p_{k-1})$, for every $k = 1, \dots, m'$;
- (c) the polygonal line $p_0 p_1 \dots p_{m'}$ is contained in P ;
- (d) the points $p_0, \dots, p_{m'}$ are not all equal.

When this property holds for some $m \in \mathbb{N}$, we simply say that P has an *orthogonal vertex connection*. We denote by $\mathcal{S}_{n,m}^* \subseteq \mathcal{P}_n$ the subset of n -gons with some orthogonal vertex connection of order m ; and set $\mathcal{S}_{n,\infty}^*$ to be the union of all $\mathcal{S}_{n,m}^*$.

Remark 5.2. A polygon P is in $\mathcal{S}_{n,\infty}^*$ iff some vertex of P has a forward orbit under the slap map which eventually hits another vertex.




FIGURE 4. Examples of polygons with orthogonal vertex connections.

Here we exclude fixed points at vertices where the polygon makes acute angles, because of condition (d).

Proposition 5.3. *The set $\mathcal{S}_{n,\infty}^*$ (resp. $\mathcal{S}_{n,m}^*$) is a countable (resp. finite) union of codimension 1, closed semialgebraic sets.*

Proof. The orthogonal projection onto an edge e with projective coordinates $(a : b : c)$ is given by

$$\pi_{(a:b:c)}(x, y) = \left(x - \frac{ax + by + c}{a^2 + b^2} a, x - \frac{ax + by + c}{a^2 + b^2} b \right). \quad (5.3)$$

Identify \mathcal{P}_n with the open subset $\Omega_n \subseteq \mathbb{P}^1 \times (\mathbb{P}^2)^{n-3} \times \mathbb{P}^1$, as explained in proposition 5.1. The set $\mathcal{S}_{n,m}^*$ is contained in the finite union of algebraic hypersurfaces defined by

$$(x_\ell, y_\ell) = \pi_{(a_{i'_m} : b_{i'_m} : c_{i'_m})} \circ \dots \circ \pi_{(a_{i_1} : b_{i_1} : c_{i_1})}(x_k, y_k) \quad (5.4)$$

where $1 \leq m' \leq m$, $(i_0 = k, i_1, \dots, i_{m'})$ is a finite sequence of indices in $\{0, 1, \dots, n-1\}$, and $v_\ell = (x_\ell, y_\ell)$ is an endpoint of e_{i_m} . This equation expresses the existence of an orthogonal vertex connection of order m between the vertices v_k and v_ℓ . Because expressions in (5.1) and (5.3) are rational functions of the coordinates $(a_i : b_i : c_i)$ of the polygon's edges, equation (5.4) is also rational in these coordinates. Reducing to a common denominator, and eliminating it, equation (5.4) becomes polynomial in the projective coordinates of $\Omega_n \subseteq \mathbb{P}^1 \times (\mathbb{P}^{n-3}) \times \mathbb{P}^1$. Notice that (5.4) is a system of two equations which reduces to single equation because the projection points lie in a line. Given distinct vertex indices i, j , and an edge index k , consider the set $\Sigma_{k, i_1, \dots, i_{m'}, \ell}$ of polygons with an orthogonal vertex connection between the vertices v_k and v_ℓ , having the prescribed itinerary. The set $\Sigma_{k, i_1, \dots, i_{m'}, \ell}$ is defined by the equation (5.4) and hence is a codimension one closed subset of the semialgebraic set \mathcal{P}_n . Thus $\mathcal{S}_{n,m}^*$ is a finite union, and $\mathcal{S}_{n,\infty}^* = \bigcup_{m \geq 1} \mathcal{S}_{n,m}^*$ a countable union, of such closed semialgebraic sets. \square

Corollary 5.4. *The set $\mathcal{P}_n \setminus \mathcal{S}_{n,m}^*$ is open and dense, whereas the set $\mathcal{P}_n \setminus \mathcal{S}_{n,\infty}^*$ is residual in \mathcal{P}_n . Both sets $\mathcal{S}_{n,m}^*$ and $\mathcal{S}_{n,\infty}^*$ have zero Lebesgue measure in \mathcal{P}_n .*

5.2. Strongly contracting generic polygonal billiards. The properties of the degenerate case $f = 0$, simpler to obtain, can be extended by Theorem 4.10 to billiards with $\lambda(f)$ very close to 0.

One can naturally reduce Φ_0 on $\{\theta = 0\}$ to a one-dimensional map called the slap map. This map is piecewise affine because the billiard is polygonal. Wherever a vertex of the polygon projects orthogonally to the interior of a side we obtain an element of the set S defined by $S_1^+ \cap \{(s, \theta) : \theta = 0\} = S \times \{0\}$. These are the pre-images of points where the slap map is not defined. On each connected open interval I_i of the domain of Φ_0 we define $\Phi_{0,i} = \Phi_0|_{I_i}$.

For any polygon we define the angles $\varphi_i \in (-\pi/2, \pi/2)$ between the straight lines supporting the sides of the polygon. If there are parallel sides we set the corresponding angle to $\varphi_i = 0$. It is simple to check that the minimum expansion rate of Φ_0 is given by

$$\alpha(\Phi_0) = \min_i |\cos \varphi_i|^{-1} \geq 1.$$

Note that $\alpha(\Phi_0) > 1$ iff there are no parallel sides.

Theorem 5.5. *For any polygon $P \in \mathcal{P}_n \setminus \mathcal{S}_{n,\infty}^*$, there is $\lambda_0 > 0$ such that if $\lambda(f) < \lambda_0$ and $f \in \mathcal{B}$, then Φ_f has a generalized hyperbolic attractor with finitely many ergodic SRB measures and dense hyperbolic periodic points.*

Proof. Notice that $\Phi_0^k(S) \cap S \neq \emptyset$ implies the existence of an orthogonal vertex connection of order k . Hence, if a polygon does not have any orthogonal vertex connection, then the corresponding slap map satisfies $\Phi_0^k(S) \cap S = \emptyset$ for every $k \geq 1$. Next observe that the number $p(S_1^+)$ depends only on the polygon P , and not on the reflection law f . Thus taking $n \gg \log p(S_1^+)/\log \alpha(\Phi_0)$, for small enough $\lambda_0 > 0$ and every $f \in \mathcal{B}$ with $\lambda(f) < \lambda_0$ one has $n > \log p(S_1^+)/\log \alpha(\Phi_f)$, and $\Phi_f^k(S_1^+) \cap S_1^+ = \emptyset$, for every $1 \leq k \leq n - 1$. The assumptions of Theorem 4.10 are met, and the theorem follows. \square

6. REGULAR POLYGONS

Throughout this section P is a regular d -gon. We assume that the sides of the polygon are normalized to a unit size. The phase space of the billiard map is thus

$$M = (0, N) \times \left(-\frac{\pi}{2}, \frac{\pi}{2}\right).$$

6.1. Phase space reduction. The study of the dynamics in regular polygonal billiards can be simplified using the group of symmetries of the polygon. In the phase space this reduction can be achieved by simply identifying all sides of the polygon. So we define an equivalence relation \sim on M by $(s_1, \theta_1) \sim (s_2, \theta_2)$ if and only if $s_1 - s_2 \in \mathbb{Z}$. Let

$$\widetilde{M} = (0, 1) \times (-\pi/2, \pi/2).$$

Clearly, the quotient space M/\sim can be identified with \widetilde{M} , which we call the *reduced phase space*. The induced billiard map ϕ_f on \widetilde{M} is the *reduced billiard map*. Notice that M is a d -fold covering of \widetilde{M} and that the reduced billiard map ϕ_f is a factor of Φ_f .

6.2. The reduced billiard map. A straightforward computation reveals that the singular set of the reduced billiard map is

$$\widetilde{N}_1^+ = \partial\widetilde{M} \cup \widetilde{S}_1^+,$$

where \widetilde{S}_1^+ is the union of the graph of the curves

$$\gamma_k(\theta) = \frac{\sin(\theta_{k-1} - \theta) \cos(\theta_k)}{\sin(\theta_{k-1} - \theta_k) \cos(\theta)}, \quad k = 2, \dots, d-1,$$

and $\theta_k = \pi/2 - k\pi/d$ which defines a partition of the interval $(-\pi/2, \pi/2)$ into d subintervals. Let,

$$A_k = \left\{ (s, \theta) \in \widetilde{M} \setminus \widetilde{N}_1^+ : \gamma_{k+1}(\theta) < s < \gamma_k(\theta) \right\}, \quad k = 1, \dots, d-1.$$

Note that $\widetilde{M} \setminus \widetilde{N}_1^+ = \cup_{k=1}^{d-1} A_k$. We then obtain the k -branch of the reduced billiard map

$$\phi_f|_{A_k}(s, \theta) = \left(\frac{\gamma_k(\theta) - s}{\gamma_k(\theta) - \gamma_{k+1}(\theta)}, f(2\theta_k - \theta) \right) \quad (6.1)$$

where f is a contracting reflection law.

Remark 6.1. If the contracting reflection law f is an odd function then ϕ_f commutes with the involution $T(s, \theta) = (1-s, -\theta)$, i.e.

$$T \circ \phi_f = \phi_f \circ T.$$

Also note that $T(A_k) = A_{d-k}$.

When $f = 0$ we obtain the one-dimensional *reduced slap map* ϕ_0 . If P is an even d -gon then $\phi_0(s) = 1-s$. Otherwise,

$$\phi_0(s) = -\frac{1}{\beta} \left(s - \frac{1}{2} \right) + \varepsilon(s),$$

where $\beta = \cos(\frac{\pi}{d})$ and $\varepsilon: [0, 1) \rightarrow \{0, 1\}$ is the function

$$\varepsilon(s) = \begin{cases} 1, & s \geq \frac{1}{2} \\ 0, & s < \frac{1}{2}. \end{cases}$$

6.3. Regular polygons with an odd number of sides.

Proposition 6.2. *For any odd number $d \geq 5$ the point $s = 1/2$ is not a pre-periodic point of the reduced slap map ϕ_0 .*

Proof. Let us write $\varepsilon_j(s) = \varepsilon(\phi_0^j(s))$ and $\delta_j(s) = 2\varepsilon_j(s) - 1$. Notice that we always have $\delta_j(s) \in \{-1, 1\}$. Then, by induction we get

$$\phi_0^n(s) = \frac{1}{2} + \frac{(-1)^n}{\beta^n} \left(s - \frac{1}{2} \right) + \frac{1}{2\beta^{n-1}} \sum_{j=0}^{n-1} (-1)^{n-j-1} \delta_j(s) \beta^j.$$

Denote by $P_n(s, \beta)$ the tail term above. More precisely, set

$$P_n(s, \beta) = \sum_{j=0}^{n-1} (-1)^{n-j-1} \delta_j(s) \beta^j.$$

Assume, by contradiction, that $\phi_0^n(1/2) = \phi_0^k(1/2)$ for some $n > k \geq 0$. Then,

$$\frac{1}{\beta^{n-1}} P_n(1/2, \beta) = \frac{1}{\beta^{k-1}} P_k(1/2, \beta),$$

which is equivalent to

$$P_n(1/2, \beta) - \beta^{n-k} P_k(1/2, \beta) = 0.$$

The last expression, hereafter denoted by $Q(\beta)$, is a polynomial in β of degree $n - 1$ with all coefficients in $\{-2, -1, 0, 1, 2\}$. Moreover, since $n > k$, $Q(0) = P_n(\frac{1}{2}, 0) = \pm 1$. Because $\tilde{\Psi}_{2d}(x)$ is the minimal polynomial of β (see the Appendix) there is a factorization $Q(x) = \tilde{\Psi}_{2d}(x) S(x)$ in $\mathbb{Q}[x]$. Since $Q(x) \in \mathbb{Z}[x]$, Gauss lemma implies there is some $c \in \mathbb{Q} \setminus \{0\}$ such that $c \tilde{\Psi}_{2d}(x) \in \mathbb{Z}[x]$ and $c^{-1} S(x) \in \mathbb{Z}[x]$. Since, by Corollary A.4, $\tilde{\Psi}_{2d}(x)$ has a unit constant term $\tilde{\Psi}_{2d}(0) = \pm 1$, it follows that $c \in \mathbb{Z}$, and hence $S(x) = c(c^{-1} S(x)) \in \mathbb{Z}[x]$. Thus, the leading coefficient of $Q(x)$ must be a multiple of the leading coefficient $\tilde{\Psi}_{2d}(x)$. But the first is ± 1 or ± 2 , while by Corollary A.4 the second has leading coefficient at least 4, which is obviously impossible. \square

Corollary 6.3. *Every odd sided regular polygon has the no orthogonal vertex connection property (see Section 5 for the definition).*

Proof. Any equilateral triangle has this property because its vertexes correspond to fixed points of the slap map. Moreover, for any odd number $d \geq 5$, if the d -sided regular polygon had a ‘orthogonal vertex connection’ this would imply that $s = 1/2$ is a pre-periodic point of the reduced slap map ϕ_0 , thus contradicting proposition 6.2. \square

Theorem 6.4. *Consider a regular polygon with an odd number of sides. There is $\lambda_0 > 0$ such that if $\lambda(f) < \lambda_0$ and $f \in \mathcal{B}$, then the billiard map Φ_f has a generalized hyperbolic attractor with finitely many ergodic SRB measures and dense hyperbolic periodic points.*

Proof. The set of singularities of the slap map includes the middle points on each of the sides of the polygon. The claim is a consequence of Theorem 4.10 and Proposition 6.2. \square

Notice that for the equilateral triangle Theorem 7.1 is an improvement of the above result.

6.4. Regular polygons with an even number of sides. The approach used to prove the existence of SRB measures for odd sided regular polygons does not work with even sided regular polygons, because these polygons have parallel sides and so the slap map is not expanding. However, this fact does not prevent the existence of hyperbolic attractors as proved for the square in Section 8 and shown by numerical experiments in [15].

In this section we show that, under certain conditions on the contracting reflection law, every orbit whose angle of incidence is sufficiently small will eventually be trapped between two parallel sides.

Consider a regular polygon with an even number $d \geq 6$ of sides. The square is considered in Section 8.

Proposition 6.5. *Let $f \in \mathcal{R}_1^k$ be an odd function such that $f' > 0$, $\lambda(f) \leq 1/2$ and $f(\delta \frac{2\pi}{d}) \leq \delta f(\frac{2\pi}{d})$ for every $0 \leq \delta \leq 1$. Then there exists a positive constant C such that every orbit of Φ_f having more than C collisions between parallel sides belongs to the basin of attraction of \mathcal{P} .*

Proof. In the reduced phase space the attractor \mathcal{P} is the interval $(0, 1)$. Moreover, since the contracting reflection law is an odd function, the set

$$B = \left\{ (s, \theta) \in \widetilde{M} : \sigma_\infty(\theta) - 1 < s < \sigma_\infty(\theta) \right\},$$

where

$$\sigma_\infty(\theta) = 1 - \cot\left(\frac{\pi}{d}\right) \sum_{n=0}^{\infty} \tan(f^n(\theta)),$$

is a trapping region, i.e., $\phi_f(B) \subset \text{int}(B)$. In fact, positive semi-orbits starting in B correspond to billiard trajectories which are trapped between any two parallel sides.

Given $\epsilon > 0$, let H_ϵ denote the horizontal strip of width ϵ around $\theta = 0$, i.e. $H_\epsilon = \left\{ (s, \theta) \in \widetilde{M} : |\theta| < \epsilon \right\}$. Also let

$$\Delta_\epsilon^\pm = A_{q \mp 1} \cap H_\epsilon \quad \text{and} \quad \Delta_\epsilon = \Delta_\epsilon^- \cup \Delta_\epsilon^+,$$


where $d = 2q$. These sets are depicted in Fig. 5. We claim that there exists an $\epsilon > 0$ such that

$$\phi_f^2(\Delta_\epsilon) \subset B. \tag{6.2}$$

Suppose that the claim holds true. It is clear that for ϵ sufficiently small $\Delta_\epsilon \subset \phi_f(A_q \setminus \text{cl}(B))$. Thus, take

$$C = \min \left\{ n \geq 1 : \phi_f^{-n}(\Delta_\epsilon) \not\subset A_q \right\}.$$

If the positive semi-orbit of $x \in \widetilde{M}$ has more than C collisions between parallel sides then, by the definition of C , there exists an $n \geq 1$

FIGURE 5. The set Δ_ϵ .

such that $\phi_f^n(x) \in \Delta_\epsilon$. It follows from the claim that $\phi_f^{n+2}(x) \in B$. Thus proving the proposition.

Therefore, it remains to prove (6.2). Since f is odd, we have by Remark 6.1 that $\phi_f \circ T = T \circ \phi_f$. Thus, it is sufficient to prove that there exists an $\epsilon > 0$ sufficiently small such that $\phi_f^2(\Delta_\epsilon^+) \subset B$.

For $q \geq 3$ the map $\phi_f|_{A_{q-1}}$ extends continuously to the boundary of A_{q-1} . Indeed, by (6.1) we have

$$\phi_f|_{A_{q-1}}(s, \theta) = \left(\cot\left(\frac{\pi}{2q}\right) \tan(g(\theta)) - s \cos(\theta) \sec(g(\theta)), f(g(\theta)) \right), \quad (6.3)$$

where $g(\theta) = \frac{\pi}{q} - \theta$. To simplify the notation, let us denote the unique extension of $\phi_f|_{A_{q-1}}$ by $\phi_{f,q-1}$.

In order to prove the inclusion $\phi_f^2(\Delta_\epsilon^+) \subset B$ for ϵ sufficiently small, we only need to trace two iterates of the point $(s, \theta) = (1, 0)$ under the continuous extension $\phi_{f,q-1}$. In fact, if $\phi_{f,q-1}(1, 0) \in \text{cl}(A_{q-1})$ and $\phi_{f,q-1}^2(1, 0) \in B$ then by continuity of $\phi_{f,q-1}$, the hypothesis $f' > 0$ and the fact that $\text{dist}(\Delta_\epsilon^+, (1, 0)) \rightarrow 0$ as $\epsilon \rightarrow 0$ we obtain the desired inclusion.

In the following we compute two iterates of the point $(s, \theta) = (1, 0)$. Taking into account (6.3), we have

$$\phi_{f,q-1}(1, 0) = (1, f(\pi/q)).$$

Since $\lambda(f) \leq 1/2$ we have that $\phi_{f,q-1}(1, 0) \in \text{cl}(A_{q-1})$. Now we show that $\phi_{f,q-1}^2(1, 0) \in B$. A simple computation reveals that

$$\phi_{f,q-1}^2(1, 0) = \left(\tan(\hat{\theta}) \left(\cot\left(\frac{\pi}{2q}\right) - \sin\left(\frac{\pi}{q}\right) \right) - \cos\left(\frac{\pi}{q}\right), f(\hat{\theta}) \right)$$

where $\hat{\theta} = g(f(\frac{\pi}{q}))$. Taking into account the definition of the set B , the inclusion $\phi_{f,q-1}^2(1, 0) \in B$ holds if and only if

$$\cot\left(\frac{\pi}{2q}\right) \sum_{n=0}^{\infty} \tan(f^{n+1}(\hat{\theta})) < 1 + \tan(\hat{\theta}) \sin\left(\frac{\pi}{q}\right) + \cos\left(\frac{\pi}{q}\right). \quad (6.4)$$

Let $\lambda_q = f(\pi/q)/(\pi/q)$. Note that $0 \leq \lambda_q \leq 1/2$. Since $f(\delta\frac{\pi}{q}) \leq \delta f(\frac{\pi}{q})$ for every $0 \leq \delta \leq 1$ we obtain the following estimate

$$f^{n+1}(\hat{\theta}) = f^{n+1}\left(\frac{\pi}{q}(1 - \lambda_q)\right) \leq \frac{\pi}{q} \lambda_q^{n+1} (1 - \lambda_q), \quad n = 0, 1, \dots$$

Thus,

$$\tan(f^{n+1}(\hat{\theta})) \leq 2 \tan\left(\frac{\pi}{2q}\right) \lambda_q^{n+1} (1 - \lambda_q),$$

by convexity of the tangent function. Using the previous estimate we get,

$$\cot\left(\frac{\pi}{2q}\right) \sum_{n=0}^{\infty} \tan(f^{n+1}(\hat{\theta})) \leq 2\lambda_q \leq 1.$$

Since $\tan(\hat{\theta}) \sin(\pi/q) + \cos(\pi/q) > 0$, inequality (6.4) holds. \square

Corollary 6.6. *Under the assumptions of Proposition 6.5, every invariant set Σ of Φ_f not intersecting the basin of attraction of \mathcal{P} is uniformly hyperbolic.*

Proof. Since Σ does not intersect the basin of attraction of \mathcal{P} then by Proposition 6.5 there exists a constant $C > 0$ such that the positive semi-orbit of every point $x \in \Sigma$ has no more than C collisions between parallel sides. Thus we can apply Proposition 3.3 and obtain the desired result. \square

Remark 6.7. The assumptions of Proposition 6.5 are satisfied for the linear contracting reflection law $f(\theta) = \sigma\theta$ where $\sigma \leq 1/2$.

Remark 6.8. In Proposition 6.5, the contracting reflection law is assumed to be an odd function. This assumption is not necessary, it only simplifies certain estimates in the proof. In fact, we can remove it provided $f(-\delta\frac{2\pi}{d}) \geq \delta f(-\frac{2\pi}{d})$ for every $0 \leq \delta \leq 1$.

7. ACUTE TRIANGLES

We now deal with the existence of SRB measures for contracting billiards in acute triangles (all the internal angles φ_i are less than $\pi/2$). The existence and uniqueness of an SRB measure for the case of the equilateral triangle and $f(\theta) = \sigma\theta$ with $\sigma < 1/3$ was first obtained in [5] by constructing a proper Markov partition.

Theorem 7.1. *For any acute triangle and $f \in \mathcal{B}$ satisfying*

$$\lambda(f) < \frac{2}{\pi} \min_i \left(\frac{\pi}{2} - \varphi_i \right),$$

the billiard map Φ_f has a generalized hyperbolic attractor with countably many ergodic SRB measures.

Proof. Let \tilde{D}_f be the invariant set in Lemma 2.3, and let $V' = V \cap \tilde{D}_f$. Hence $\Phi_f(S_1^+) \subset V'$. Since S_1^+ intersects V at $\theta = \pi/2 - \varphi_i$, it is not difficult to see that if $\lambda(f) < 2 \min_i (\pi/2 - \varphi_i) / \pi$, then $S_1^+ \cap V' = \emptyset$ and $\Phi_f(V') \subset V'$. Therefore, $\Phi_f^k(S_1^+) \cap S_1^+ = \emptyset$ for every $k \geq 1$. The wanted conclusion now follows from Theorem 4.9. \square

8. RECTANGLES

Throughout this section, the polygon P is a rectangle. Recall that \mathcal{P} denotes the set of periodic points of period two of Φ , and let $B(\mathcal{P})$ be the basin of attraction of \mathcal{P} . In the following, we show that although $\mathcal{P} \neq \emptyset$, the invariant set $D \setminus B(\mathcal{P})$ may still contain non-trivial hyperbolic sets. We also give sufficient conditions for $B(\mathcal{P}) = M \setminus N_\infty^+$.

We start by showing that $D \setminus B(\mathcal{P})$ is always hyperbolic.

Proposition 8.1. *The set $D \setminus B(\mathcal{P})$ is hyperbolic.*

Proof. A sequence of consecutive collisions $\{(s_i, \theta_i)\}_{i=0}^m$ with $m \geq 1$ is called a *block* if its first m collisions occur at two parallel sides of P , whereas its last collision occurs at a side adjacent to the other two; when $m = 1$, a block consists of two consecutive collisions at adjacent sides of the rectangle. It is easy to see that $|\theta_{m-1}| \leq \lambda^{m-1} |\theta_0|$, and using (2.1), that

$$|\bar{\theta}_m - \theta_{m-1}| = \frac{\pi}{2}. \quad (8.1)$$

Since $\rho \geq 1$, it follows that $\Lambda_m(s_1, \theta_1) \geq \rho(\bar{\theta}_m)$. Next, we want to estimate $\rho(\bar{\theta}_m)$ from below. Let

$$m_\lambda = \frac{2}{\log \lambda} \log \left(\frac{2}{\pi} \cos \frac{\pi \lambda}{2} \right) \quad \text{and} \quad b_\lambda = r \left(\frac{\pi}{2} (1 - \lambda) \right).$$

We consider separately the cases: $m \leq m_\lambda$ and $m > m_\lambda$. First, suppose that $m \leq m_\lambda$. By Lemma 2.3, we know that $|\theta_{m-1}| < \pi \lambda / 2$. Using (8.1), we obtain $|\bar{\theta}_m| > \pi(1 - \lambda)/2$ so that

$$\Lambda_m(s_1, \theta_1) \geq b_\lambda \geq b_\lambda^{\frac{m}{m_\lambda}}. \quad (8.2)$$

Now, suppose that $m > m_\lambda$. Using the bounds $|\theta_0| < \lambda \pi / 2$ and $|\theta_m| < \lambda \pi / 2$ provided by Lemma 2.3, we get

$$\cos(|\bar{\theta}_m|) \leq \sin(\lambda^{m-1} |\theta_0|) < \sin \frac{\pi \lambda^m}{2} \leq \frac{\pi \lambda^m}{2}, \quad (8.3)$$

and

$$\cos(|\theta_m|) \geq \cos \frac{\pi \lambda}{2}. \quad (8.4)$$

Since $m > m_\lambda$, it follows from the definition of m_λ that

$$\pi\lambda^{m/2} \leq 2 \cos \frac{\pi\lambda}{2}.$$

This fact, together with (8.4) and (8.3), implies that

$$\begin{aligned} \Lambda_m(s_1, \theta_1) &\geq \rho(\bar{\theta}_m) > \frac{2}{\pi\lambda^m} \cos \frac{\pi\lambda}{2} \\ &\geq \lambda^{-\frac{m}{2}} \frac{2}{\pi\lambda^{\frac{m}{2}}} \cos \frac{\pi\lambda}{2} \geq \lambda^{-\frac{m}{2}}. \end{aligned} \quad (8.5)$$

Let $\mu = \min\{b_\lambda^{1/m_\lambda}, \lambda^{-1/2}\} > 1$. Combining (8.2) and (8.5) together, we conclude that

$$\Lambda_m(s_1, \theta_1) \geq \mu^m. \quad (8.6)$$

Suppose that $(s_0, \theta_0) \in D \setminus B(\mathcal{P})$. Then for $n > 0$ sufficiently large, the sequence of collisions $\{(s_i, \theta_i)\}_{i=0}^n$ can be decomposed into $k(n) \geq 1$ consecutive blocks followed by $j(n) \geq 0$ consecutive collisions with two parallel sides of the rectangle. Let $L(s_0, \theta_0; n)$ be positive integer such that $x_{L(s_0, \theta_0; n)}$ is the last collision of the $k(n)$ th block. Hence

$$L(s_0, \theta_0; L(s_0, \theta_0; n)) = L(s_0, \theta_0; n) \leq n.$$

Since $(s_0, \theta_0) \notin B(\mathcal{P})$, we have $L(s_0, \theta_0; n) \rightarrow +\infty$ as $n \rightarrow +\infty$, and so

$$\limsup_{n \rightarrow +\infty} \frac{L(s_0, \theta_0; n)}{n} = 1.$$

Let $A = \cos(\pi\lambda/2)$. From (8.6), it follows that

$$\alpha_n(s_0, \theta_0) \geq A\mu^{L(s_0, \theta_0; n)}$$

for n sufficiently large. But $\|D\Phi^n(s_0, \theta_0)\| \geq \alpha_n(s_0, \theta_0)$ so that

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log \|D\Phi^n(s, \theta)\| \geq \log \mu > 0.$$

This and Proposition 3.1 imply that $D \setminus B(\mathcal{P})$ is hyperbolic. \square

Remark 8.2. Suppose that $j(n)$ in the proof of Proposition 8.1 is bounded from above by some constant J for every n and every $(s_0, \theta_0) \in D \setminus B(\mathcal{P})$. Then $D \setminus B(\mathcal{P})$ is uniformly hyperbolic by Proposition 3.3. We can also obtain the same conclusion, in a more direct way, by arguing as follows. If such a J exists, then $L(s_0, \theta_0) \geq n - J$. The notation here is as in the proof of Proposition 3.3. We then obtain,

$$\alpha_n(s_0, \theta_0) \geq A\mu^{L(s_0, \theta_0; n)} \geq \frac{A}{\mu^J} \mu^n,$$

which means that $D \setminus B(\mathcal{P})$ is uniformly hyperbolic.

Remark 8.3. Since the estimate (8.6) relies only on the assumption that a rectangle has three consecutive sides with internal angles equal to $\pi/2$, one can easily extend Proposition 8.1 to polygons that are finite unions of rectangles with sides parallel to two orthogonal axes.

Remark 8.4. It is not difficult to see that $B(\mathcal{P}) = M \setminus N_\infty^+$ for every $f \in \mathcal{R}_1^k$ with $f' < 0$. This kind of reflection law is the contracting version of $f(\theta) = -\theta$, the reflection law of Andreev billiards, which are models for the electronic conduction in a metal connected to a super-conductor [24].

We now give some sufficient conditions for the existence of a forward invariant set of positive ν -measure containing a uniformly hyperbolic set. In view of Remark 8.4, we may assume that $f' > 0$. Moreover, by rescaling P , we may assume without loss of generality that the sides of P have length 1 and $0 < h \leq 1$. Fix a pair of parallel sides of P . In the following, the expression ‘long sides’ (resp. ‘long side’) of P has to be understood as the chosen pair (resp. a side from the chosen pair) of parallel sides of P whenever $h = 1$.

Define

$$f_1(\theta) = -f(\theta) \quad \text{for } \theta \in [\pi/2, \pi/2],$$

and

$$f_2(\theta) = f\left(\operatorname{sgn}(\theta)\frac{\pi}{2} - \theta\right) \quad \text{for } \theta \in [-\pi/2, 0) \cup (0, \pi/2].$$

To explain the geometrical meaning of f_1 and f_2 , we observe that if (s_0, θ_0) and (s_1, θ_1) are two consecutive collisions, then $\theta_1 = f_1(\theta_0)$ when the two collisions occur at parallel sides of P , and $\theta_1 = f_2(\theta_0)$ when the two collisions occur at adjacent sides of P .

From the properties of f , one can easily deduce that i) f_2 is strictly decreasing, ii) the restriction of f_2 to each interval $(-\pi/2, 0)$ and $(0, \pi/2)$ is a strict contraction with Lipschitz constant less than $\lambda(f)$, and iii) f_2 has two fixed points $\theta^- < 0 < \theta^+$.

Definition 8.1. Let T be the subset of $M \setminus N_\infty^+$ consisting of elements whose positive semi-trajectory is trapped between two parallel sides of P .

It is easy to see that T is a forward invariant set containing \mathcal{P} .

Definition 8.2. Let $\hat{M} = M \setminus (T \cup N_\infty^+)$.

The set \hat{M} consists of elements with infinite positive semi-trajectory having at least a pair of consecutive collisions at adjacent sides of P . It can be shown that \hat{M} is open up to a set of zero ν -measure.

Definition 8.3. The sequence $\Phi(x_0), \dots, \Phi^n(x_0)$ of $n \geq 2$ collisions with $x_0 \in \hat{M}$ is called *adjacent* if $\Phi^i(x_0)$ and $\Phi^{i+1}(x_0)$ hit adjacent sides of P for every $i = 1, \dots, n-1$.

Definition 8.4. The sequence $\Phi(x_0), \dots, \Phi^{n-1}(x_0)$ of $n \geq 1$ collisions hitting two parallel sides of P with $x_0 \in \hat{M}$ is called *complete* if x_0 and x_1 are adjacent, and x_{n-1} and x_n are adjacent. The number n is called the *length* of the sequence.

Let $\theta_*^- < 0 < \theta_*^+$ be the two solutions of

$$\mathcal{F}(\theta) := \sum_{n=0}^{\infty} \tan(|f_1^n(\theta)|) = \frac{1}{h}, \quad \theta \in (-\pi/2, \pi/2).$$

If the endpoints of a long side of P have arclength parameters $s = 0$ and $s = 1$, then the billiard particle leaving the endpoint $s = 0$ (resp. $s = 1$) with an angle θ_*^+ (resp. θ_*^-) bounces between the long sides of P forever, converging to the short side of P that has an endpoint at $s = 1$ (resp. $s = 0$).

Lemma 8.5. *If $(s_0, \theta_0) \in M \setminus N_{\infty}^+$ is the first element of a complete sequence of collisions between the long sides of P and*

$$f_1(\theta_*^+) \leq \theta_0 \leq f_1(\theta_*^-), \quad (8.7)$$

then $(s_0, \theta_0) \in T$.

Proof. From the monotonicity of f and Condition (8.7), it follows that $\mathcal{F}(\theta_0) < 1/h$. This implies immediately the wanted conclusion. \square

Lemma 8.6. *If $(s_0, \theta_0) \in M \setminus N_{\infty}^+$ is the first element of a complete sequence of collisions between parallel sides of P and*

$$\theta_0 < f_1(\theta_*^+) \quad \text{or} \quad f_1(\theta_*^-) < \theta_0, \quad (8.8)$$

then (s_m, θ_m) and (s_{m+1}, θ_{m+1}) are adjacent for some $m \geq 0$.

Proof. Using the monotonicity of f and Condition (8.8), we obtain $\mathcal{F}(\theta_0) > 1/h > h$, which proves the proposition. \square

Let

$$\tilde{\theta} := \min \left\{ f_2 \left(f \left(\frac{\pi}{2} \right) \right), -f_2 \left(f \left(-\frac{\pi}{2} \right) \right) \right\}.$$

Lemma 8.7. *Let $(s_0, \theta_0) \in \tilde{D}$. If (s_0, θ_0) and (s_1, θ_1) are adjacent collisions, then $|\theta_1| \geq \tilde{\theta}$.*

Proof. We have $|\theta_1| = |f_2(f(\bar{\theta}_0))|$, because $\theta_1 = f_2(\theta_0)$ and $\theta_0 = f(\bar{\theta}_0)$ with $\bar{\theta}_0 \in (-\pi/2, \pi/2)$. But $(f_2 \circ f)' < 0$, since $f' > 0$. Thus, the minimum of $|f_2 \circ f|$ is attained at $-\pi/2$ or $\pi/2$, and therefore it is equal to $\tilde{\theta}$. \square

Proposition 8.8. *Suppose that $f' > 0$ and*

$$\tilde{\theta} > \max\{-\theta_*^-, \theta_*^+\}. \quad (8.9)$$

Then there exists $m > 0$ such that for every $(s_0, \theta_0) \in \hat{M}$, the positive semi-orbit $\{(s_n, \theta_n)\}_{n \geq 0}$ contains infinitely many complete sequences, and each complete sequence contained in $\{(s_n, \theta_n)\}_{n \geq 0}$ has length less or equal than m .

Proof. We can assume without loss of generality that $s = 0$ and $s = 1$ correspond to the endpoints of a long side of P , and $s = 1$ and $s = 1 + h$ correspond to the endpoints of the adjacent short side. By (8.9), we can apply Lemma 8.6 to the four collisions $(0, \tilde{\theta}), (1, -\tilde{\theta}), (1, \tilde{\theta}), (1, -\tilde{\theta})$ with the angle measured with respect to the normal to the long side for the first two collisions and with respect to the normal of the short side for the last two collisions. We conclude that there exists $n > 0$ such that the first pair of adjacent collisions in the semi-orbit of each of $(0, \tilde{\theta}), (1, -\tilde{\theta}), (1, \tilde{\theta}), (1, -\tilde{\theta})$ comes after a sequence of no more than $n > 0$ consecutive collisions between two parallel sides of P .

Now, let $x_0 \in \hat{M}$. From the definition of \hat{M} , there exists $m > 0$ such that x_m and x_{m+1} are adjacent collisions. Hence $\theta_{m+1} = f_2(\theta_m)$, and from Lemma 8.7, it follows that $|\theta_{m+1}| \geq \tilde{\theta}$. The conclusion obtained in the previous paragraph implies that we can find $1 \leq k \leq n$ such that the collisions x_{m+k} and x_{m+k+1} are adjacent. To complete the proof, we just need to observe that n depends only on $\tilde{\theta}$. \square

Corollary 8.9. *Under the hypotheses of Proposition 8.8, we have*

- (1) \hat{M} is a forward invariant set,
- (2) every invariant subset of \hat{M} is uniformly hyperbolic,
- (3) $B(\mathcal{P}) = T$.

Proof. Proposition 8.8 implies that \hat{M} is forward invariant and has Property (A). This gives Part (1), and by using Proposition 3.3, Part (2) as well. Part (3) follows from the definition of \hat{M} . \square

Remark 8.10. From Corollary 8.9, it follows that \hat{M} contains a hyperbolic attractor \mathcal{H} . This gives us the following dynamical picture: up to a set of zero ν -measure, M is the union (mod 0) of two positive ν -measure sets, the basin of attraction of \mathcal{P} and the basin of attraction of \mathcal{H} . Although Theorem 4.4 is formulated for polygons with $\mathcal{P} = \emptyset$, it is not difficult to see that we can apply it to \mathcal{H} . The challenging problem here is to prove that for a rectangle (satisfying the hypotheses of the corollary) the hypothesis $p(S_m^+) < \alpha(\Phi^m)$ for some $m > 0$ of Theorem 4.4 is satisfied.

Remark 8.11. By modifying properly the definition of $\tilde{\theta}$ and Condition 8.9, it is possible to extend Proposition 8.8 to a larger class of polygons with parallel sides.

Finally, we give sufficient conditions for having $B(\mathcal{P}) = M \setminus N_\infty^+$. Recall that θ^- and θ^+ are the fixed points of f_2 .

Proposition 8.12. *Suppose that $f' > 0$ and*

$$h \leq \tan \tilde{\theta} \tag{8.10}$$

$$f_1(\theta_*^+) < \theta^- \quad \text{and} \quad \theta^+ < f_1(\theta_*^-). \tag{8.11}$$

Then $B(\mathcal{P}) = M \setminus N_\infty^+$.

Proof. Let $x_0 \in M \setminus N_\infty^+$. As usual, we write $x_n = (s_n, \theta_n) = \Phi^n(x_0)$. If $\theta_0 = 0$, then $x_0 \in \mathcal{P}$, and there is nothing to prove. Hence, we may assume that $\theta_0 \neq 0$. In this case, arguing by contradiction, we show that there exists $n > 0$ such that $x_n \in T$.

Suppose that $x_n \notin T$ for every $n \geq 0$. Then two cases can occur: either there exists $m > 0$ such that all collisions x_m, x_{m+1}, \dots are adjacent, or there exist two sequences $\{m_k\}$ and $\{n_k\}$ of positive integers such that $0 \leq m_k < n_k < m_{k+1}$ and x_{m_k}, \dots, x_{n_k} is a complete sequence of collisions between the long sides of P for every $k \geq 0$.

In the first case, we have

$$\theta_n = f_2^{n-m}(\theta_m) \quad \text{for } n \geq m.$$

By the properties of f_2 , it follows that either $\theta_n \rightarrow \theta^-$ or $\theta_n \rightarrow \theta^+$ as $n \rightarrow +\infty$. Condition (8.11) guarantees the existence of $n > m$ such that x_n is a collision at one of the long sides of P , and $f_1(\theta_+^*) < \theta_n < f_1(\theta_-^*)$. Lemma 8.5 implies that $x_n \in T$, obtaining a contradiction.

We now consider the second case. Using Lemma 8.7, we see that Condition (8.10) implies that x_{n_k+1} and x_{n_k+2} are adjacent collisions. The same is true, as a direct consequence of the definition of m_k and n_k , for the sequences x_{n_k}, x_{n_k+1} and $x_{n_k+2}, \dots, x_{m_{k+1}}$ (possibly with $n_k + 2 = m_{k+1}$). Hence

$$\theta_{m_{k+1}} = f_2^{m_{k+1}-n_k}(\theta_{n_k}).$$

If $m_{k+1} - n_k$ is odd, then from the properties of f_2 , we deduce that either $0 < \theta_{m_{k+1}} < \theta^-$ or $0 < \theta_{m_{k+1}} < \theta^+$. This together with Condition (8.11) and Lemma 8.5 gives the contradiction $x_{m_{k+1}} \in T$. Now, suppose that $m_{k+1} - n_k$ is even. The assumption $x_{m_k} \notin T$ implies that $\theta_{m_{k+1}} < f_1(\theta_+^*)$ or $f_1(\theta_-^*) < \theta_{m_{k+1}}$. Using Condition (8.11), we see that $\theta_{m_{k+1}} < \theta^-$ or $\theta^+ < \theta_{m_{k+1}}$. This and the fact that f_2^2 is increasing imply that

$$|\theta_{m_{k+1}}| \leq |\theta_{n_k}|.$$

Since $\theta_{n_k} = f_1^{n_k-m_k}(\theta_{m_k})$, we have

$$|\theta_{m_{k+1}}| \leq |f_1^{n_k-m_k}(\theta_{m_k})| \leq \lambda^{n_k-m_k} |\theta_{m_k}|.$$

From this, it follows that $|\theta_{m_k}| \rightarrow 0$ as $k \rightarrow +\infty$. Thus there exists $k > 0$ such that $f_1(\theta_+^*) < \theta_{m_k} < f_1(\theta_-^*)$, and again using Lemma 8.5, we obtain the contradiction $x_{m_k} \in T$. This completes the proof. \square

Remark 8.13. When f is odd, the hypotheses of Propositions 8.8 and 8.12 simplify significantly. Indeed, in this case, we have $\theta_*^- = -\theta_*^+$, $\theta^- = -\theta^+$ and $\theta = f(\pi/2 - f(\pi/2))$.

Remark 8.14. If $f(\theta) = \sigma\theta$, then it is easy to show that Condition 8.9 is satisfied for every $0 < \sigma < 1$ and every $0 < h \leq 1$ such that $2/\pi < \sigma h$ (see also [15, Proposition V.9] for the square). Moreover, given $0 < \sigma < 1$, Conditions (8.10) and (8.11) are satisfied for h sufficiently small.

APPENDIX A. PROOFS OF SOME TECHNICAL RESULTS

Proof of Proposition 2.1. Let $B_k = [\tilde{s}_k, \tilde{s}_{k+1}] \times (-\pi/2, \pi/2)$ for $1 \leq k \leq n$, and let $\gamma_{jk} = q_1^{-1}(C_j) \cap B_k$ with $1 \leq j < k$ or $k+1 < j \leq n$. Then we have

$$S_1^+ = \left(\bigcup_{\substack{k=1 \\ 1 \leq j \leq k-1}}^n \overline{\gamma_{jk}} \right) \cup \left(\bigcup_{\substack{k=1 \\ k+2 \leq j \leq n}}^n \overline{\gamma_{jk}} \right).$$

We show that the sets $\overline{\gamma_{jk}}$ are indeed the curves $\Gamma_1, \dots, \Gamma_m$. Fix j and k . We can always find a Cartesian coordinate system (X, Y) of \mathbb{R}^2 such that the side L_k of P lies on the X -axis, and has the same orientation as the X -axis. If $(s, \theta) \in \gamma_{jk}$, then

$$\theta = \varphi_{jk}(s) := \arctan \left(\frac{\bar{s} - s}{l} \right),$$

where l is the distance between C_j and L_k , and \bar{s} is the abscissa of the vertex C_j . It is easy to see that γ_{jk} is the graph of φ_{jk} over an open interval $I_{jk} \subsetneq (\tilde{s}_k, \tilde{s}_{k+1})$ if there exist $x \in \partial\gamma_{jk} \setminus V$ and a vertex C_i with $i \neq j$ such that $q_1(x) = C_i$. Otherwise, γ_{jk} is the graph of φ_{jk} over the entire interval $(\tilde{s}_k, \tilde{s}_{k+1})$. The function φ_{jk} admits a unique smooth extension up to the boundary of I_{jk} . If we denote by $\bar{\varphi}_{jk}$ such an extension, then $\overline{\gamma_{jk}} = \text{graph}(\bar{\varphi}_{jk})$. Moreover,

$$\bar{\varphi}'_{jk}(s) = -\frac{l}{l^2 + (\bar{s} - s)^2} < 0 \quad \text{for } s \in \overline{I_{jk}}. \quad (\text{A.1})$$

It is not difficult to see from (A.1) that if $\bar{\varphi}_{ik}(s) = \bar{\varphi}_{jk}(s)$ for some $s \in [\tilde{s}_k, \tilde{s}_{k+1}]$ and some $i \neq j$, then $\bar{\varphi}'_{ik}(s) \neq \bar{\varphi}'_{jk}(s)$. This conclusion and previous observations prove Parts (1)-(3) of the proposition.

Part (4) of the proposition is proved by contradiction. Suppose that there exists $x \in \gamma_{ik} \cap \gamma_{jk}$ for some $i \neq j$. Then we would have simultaneously $q_1(x) = C_i$ and $q_1(x) = C_j$, implying that $i = j$. This completes the proof. \square

Proof of Proposition 2.2. First of all, recall that S_1^+ and S_1^- are diffeomorphic. Since $DF(s, \theta) = \text{diag}(1, -f'(\theta))$, the curves forming S_1^- are strictly increasing if $f' > 0$ and strictly decreasing if $f' < 0$. \square

Proof of Proposition 2.4. The Jacobian $J(s_0, \theta_0)$ of Φ_f at $(s_0, \theta_0) \in D_f$ is given by

$$J(s_0, \theta_0) = \left| \frac{\cos \theta_0}{\cos \bar{\theta}_1} \right| \cdot |f'(\bar{\theta}_1)|, \quad (\text{A.2})$$

where $(s_1, \theta_1) = \Phi_f(s_0, \theta_0)$. By Formula (2.1), we have

$$\bar{\theta}_1 = \pi - \delta(L_0, L_1) - \theta_0, \quad (\text{A.3})$$

where L_0 and L_1 are the sides of P containing $q(s_0, \theta_0)$ and $q(s_1, \theta_1)$, respectively. The hypothesis of the proposition implies that there exists $a > 0$ such that

$$\cos(\pi - \delta(L_0, L_1)) \geq a \quad \text{for } (s_0, \theta_0) \in D_f. \quad (\text{A.4})$$

From (A.2)-(A.4) and the fact that $|\theta_0| \leq \lambda(f)\pi/2$, it follows that

$$J(s_0, \theta_0) \leq \frac{\lambda(f)}{|\cos(\pi - \delta - \theta_0)|} \xrightarrow{\lambda(f) \rightarrow 0^+} 0$$

uniformly in $(s_0, \theta_0) \in D_f$. Hence, there exists $\bar{J} < 1$ such that for $\lambda(f)$ sufficiently small, we have $J(s_0, \theta_0) \leq \bar{J}$ for every $(s_0, \theta_0) \in D_f$. This easily implies the conclusion of the proposition. \square

Proof of Proposition 2.5. The proposition follows immediately from the fact that $J(s_0, \theta_0) < 1$ for every $(s_0, \theta_0) \in D_f$, which in turn is a direct consequence of (A.2) and the hypothesis of the proposition. \square

We now establish some facts about Chebyshev polynomials that are used in Section 6 to analyze the slap map of odd sided regular polygons. The slopes of these maps are trigonometric algebraic numbers having Chebyshev polynomials as their minimal polynomials. We use the following lemma due to Gauss [30, Lemma 3.17].

Lemma A.1. *Given a polynomial $F(x) \in \mathbb{Z}[x]$, if $F(x)$ factors as $P(x)Q(x)$ in $\mathbb{Q}[x]$ it also factors as $\overline{P}(x)\overline{Q}(x)$ in $\mathbb{Z}[x]$. More precisely there is some constant $c \in \mathbb{Q} \setminus \{0\}$ such that $\overline{P}(x) = cP(x) \in \mathbb{Z}[x]$ and $\overline{Q}(x) = c^{-1}Q(x) \in \mathbb{Z}[x]$. Hence, given a polynomial in $\mathbb{Z}[x]$, it is irreducible over $\mathbb{Z}[x]$ iff it is irreducible over $\mathbb{Q}[x]$.*

The Chebyshev polynomials of the first kind are defined recursively by

$$\begin{aligned} T_0(x) &= 1, \quad T_1(x) = x, \\ T_{n+1}(x) &= 2xT_n(x) - T_{n-1}(x), \end{aligned}$$

from where it follows easily, by induction, that

$$T_n(\cos(\theta)) = \cos(n\theta). \quad (\text{A.5})$$

Chebyshev polynomials are related in [32] with the minimal polynomials $\Psi_n(x)$ of the trigonometric algebraic numbers of the form $\beta_n = \cos(\frac{2\pi}{n})$, which are real parts of unity roots. The minimal polynomial $\Psi_n(x)$ is considered there as a monic polynomial with rational coefficients, i.e., $\Psi_n(x) \in \mathbb{Q}[x]$. The relation established is the following:

Theorem A.2 ([32]). *For any integer $n \geq 1$,*

- (a) *if $n = 2s + 1$, then $T_{s+1}(x) - T_s(x) = 2^s \prod_{d|n} \Psi_d(x)$;*
- (b) *if $n = 2s$, then $T_{s+1}(x) - T_{s-1}(x) = 2^s \prod_{d|n} \Psi_d(x)$.*

This theorem allows a recursive calculation of the polynomials $\Psi_n(x)$. One obtains for instance $\Psi_1(x) = x - 1$, $\Psi_2(x) = x + 1$, $\Psi_3(x) = x + \frac{1}{2}$, $\Psi_4(x) = x$ and $\Psi_5(x) = x^2 + \frac{1}{2}x - \frac{1}{4}$. Define now $\tilde{\Psi}_1(x) = \Psi_1(x) = x - 1$, and $\tilde{\Psi}_n(x) = 2^{\deg(\Psi_n)} \Psi_n(x)$ for every $n \geq 2$. Denote by $\phi(n)$ the Euler function, which counts the number of integers co-prime with n in the range $[1, n]$.

Corollary A.3. *For $n > 2$, $\tilde{\Psi}_n(x)$ is a polynomial in $\mathbb{Z}[x]$ of degree $\phi(n)/2$. Moreover,*

- (a) *if $n = 2s + 1$, then $T_{s+1}(x) - T_s(x) = \prod_{d|n} \tilde{\Psi}_d(x)$, and*
- (b) *if $n = 2s$, then $T_{s+1}(x) - T_{s-1}(x) = \prod_{d|n} \tilde{\Psi}_d(x)$.*

Proof. The degree of the polynomial $\Psi_n(x)$, or $\tilde{\Psi}_n(x)$, was computed in [19] to be $\phi(n)/2$. The proof goes by induction in n , using Theorem A.2 and Gauss' lemma. Notice that $\deg(T_s) = s$. \square

From the previous computations we get, for instance, $\tilde{\Psi}_1(x) = x - 1$, $\tilde{\Psi}_2(x) = 2x + 2$, $\tilde{\Psi}_3(x) = 2x + 1$, $\tilde{\Psi}_4(x) = 2x$ and $\tilde{\Psi}_5(x) = 4x^2 + 2x - 1$.

Corollary A.4. *For any $n \geq 5$, the minimal polynomial $\tilde{\Psi}_n$ has leading coefficient 2^r , where $r = \phi(n)/2$, and unit constant term $\tilde{\Psi}_n(0) = \pm 1$. Moreover, $r = \phi(n)/2 \geq 2$ when $n \geq 7$.*

Proof. By recursion, $T_n(0) = 0$ if n is odd, and $T_n(0) = \pm 1$ if n is even. Hence, if $n = 2s + 1$, then $T_{s+1}(0) - T_s(0) = \pm 1$, if $n = 2s$ with s odd, then $T_{s+1}(0) - T_{s-1}(0) = \pm 2$, while if $n = 2s$ with s even, then $T_{s+1}(0) - T_{s-1}(0) = 0$. Note now that $\tilde{\Psi}_2(0) = 2$ and $\tilde{\Psi}_4(0) = 0$. By induction we can prove that $\tilde{\Psi}_n(0) = \pm 1$ for every $n \geq 5$. Finally, we have $\phi(6) = 2$, but using that $\phi(n)$ is multiplicative it follows that $2r = \phi(n) \geq 4$ for every $n \geq 7$. \square

ACKNOWLEDGEMENTS

The authors were supported by Fundação para a Ciência e a Tecnologia through the Program POCI 2010 and the Project “Randomness in Deterministic Dynamical Systems and Applications” (PTDC-MAT-105448-2008). G. Del Magno wish to express their thanks to M. Lenci and R. Markarian for stimulating conversations.

REFERENCES

- [1] V. S. Afraïmovich and Ya. B. Pesin, Dimension of Lorenz type attractors, in *Mathematical physics reviews*, Vol. 6, 169–241, Soviet Sci. Rev. Sect. C Math. Phys. Rev., 6 Harwood Academic Publ., Chur.
- [2] V. S. Afraïmovich, N. I. Chernov and E. A. Sataev, Statistical properties of 2-D generalized hyperbolic attractors, *Chaos* **5** (1995), no. 1, 238–252.

- [3] E. G. Altmann, G. Del Magno and M. Hentschel, Non-Hamiltonian dynamics in optical microcavities resulting from wave-inspired corrections to geometric optics, *Europhys. Lett. EPL* **84** (2008), no. 1, Art. 10008, 6 pp.
- [4] A. Arroyo, R. Markarian and D. P. Sanders, Bifurcations of periodic and chaotic attractors in pinball billiards with focusing boundaries, *Nonlinearity* **22** (2009), no. 7, 1499–1522.
- [5] A. Arroyo, R. Markarian, and D. P. Sanders, Structure and evolution of strange attractors in non-elastic triangular billiards, *Chaos* **22**, 2012, 026107.
- [6] V. M. Belykh, *Qualitative methods of the theory of nonlinear oscillations in point systems*, Gorki University Press, 1980.
- [7] L. A. Bunimovič, The ergodic properties of certain billiards, *Funkcional. Anal. i Priložen.* **8** (1974), no. 3, 73–74.
- [8] L. A. Bunimovich, On absolutely focusing mirrors, in *Ergodic theory and related topics, III (Güstrow, 1990)*, 62–82, Lecture Notes in Math., 1514 Springer, Berlin.
- [9] L. Bunimovich, Billiards and other hyperbolic systems, in *Encyclopedia of Mathematical Sciences*, 100, Springer-Verlag, New York, 2000, 192–233.
- [10] N. Chernov and H.-K. Zhang, On statistical properties of hyperbolic systems with singularities, *J. Stat. Phys.* **136** (2009), no. 4, 615–642.
- [11] N. Chernov and R. Markarian, *Chaotic billiards*, Mathematical Surveys and Monographs, 127, Amer. Math. Soc., Providence, RI, 2006.
- [12] N. I. Chernov, A. Korepanov and N. Simányi, Stable regimes for hard disks in a channel with twisting walls, *Chaos* **22** (2012), 026105.
- [13] V. J. Donnay, Using integrability to produce chaos: billiards with positive entropy, *Comm. Math. Phys.* **141** (1991), no. 2, 225–257.
- [14] E. Gutkin, Billiards in polygons: survey of recent results, *J. Statist. Phys.* **83** (1996), no. 1-2, 7–26.
- [15] G. Del Magno, J. Lopes Dias, P. Duarte, J. P. Gaivão and D. Pinheiro, Chaos in the square billiard with a modified reflection law, *Chaos* **22**, (2012), 026106.
- [16] G. Del Magno, J. Lopes Dias, P. Duarte, J. P. Gaivão and D. Pinheiro, Ergodicity of contractive polygonal billiards, in preparation.
- [17] A. Katok, The growth rate for the number of singular and periodic orbits for a polygonal billiard, *Comm. Math. Phys.* **111** (1987), no. 1, 151–160.
- [18] A. Katok et al., *Invariant manifolds, entropy and billiards; smooth maps with singularities*, Lecture Notes in Mathematics, 1222, Springer, Berlin, 1986.
- [19] D. H. Lehmer, Questions, Discussions, and Notes: A Note on Trigonometric Algebraic Numbers, *Amer. Math. Monthly* **40** (1933), no. 3, 165–166.
- [20] R. Lozi, Un attracteur étrange du type attracteur de Hénon, *J. Phys. C* **39** Coll. C5, (1978), 9–10.
- [21] R. Markarian, E. J. Pujals and M. Sambarino, Pinball billiards with dominated splitting, *Ergodic Theory Dynam. Systems* **30** (2010), no. 6, 1757–1786.
- [22] R. Markarian, Billiards with Pesin region of measure one, *Comm. Math. Phys.* **118** (1988), no. 1, 87–97.
- [23] Ya. B. Pesin, Dynamical systems with generalized hyperbolic attractors: hyperbolic, ergodic and topological properties, *Ergodic Theory Dynam. Systems* **12** (1992), no. 1, 123–151.
- [24] K. Richter, *Semiclassical Theory of Mesoscopic Quantum Systems*, Springer Tracts in Modern Physics, 161, Springer-Verlag, 1999.
- [25] E. A. Sataev, *Uspekhi Mat. Nauk* **47** (1992), no. 1(283), 147–202, 240; translation in *Russian Math. Surveys* **47** (1992), no. 1, 191–251.
- [26] Ja. G. Sinaï, Dynamical systems with elastic reflections. Ergodic properties of dispersing billiards, *Uspekhi Mat. Nauk* **25** (1970), no. 2 (152), 141–192.

- [27] J. Schmeling and S. Troubetzkoy, Dimension and invertibility of hyperbolic endomorphisms with singularities, *Ergodic Theory Dynam. Systems* **18** (1998), no. 5, 1257–1282.
- [28] *Hard ball systems and the Lorentz gas*, Encyclopaedia of Mathematical Sciences, 101, Springer, Berlin, 2000.
- [29] S. Tabachnikov, Billiards, *Panor. Synth.* No. 1 (1995), vi+142 pp.
- [30] I. Stewart, *Galois Theory*, third edition, Chapman & Hall/CRC Mathematics, Chapman & Hall/CRC, Boca Raton, FL, 2004.
- [31] J. Smillie, The dynamics of billiard flows in rational polygons, *Encyclopedia of Mathematical Sciences*, 100, Springer-Verlag, New York, 2000, 360–382.
- [32] W. Watkins and J. Zeitlin, The minimal polynomial of $\cos(2\pi/n)$, *Amer. Math. Monthly* **100** (1993), no. 5, 471–474.
- [33] M. Wojtkowski, Principles for the design of billiards with nonvanishing Lyapunov exponents, *Comm. Math. Phys.* **105** (1986), no. 3, 391–414.
- [34] L.-S. Young, Bowen-Ruelle measures for certain piecewise hyperbolic maps, *Trans. Amer. Math. Soc.* **287** (1985), no. 1, 41–48.
- [35] L.S. Young, Statistical properties of dynamical systems, *Annals Math.* **147** (1998), no. 3, 585–650.
- [36] L.-S. Young, What are SRB measures, and which dynamical systems have them?, *J. Statist. Phys.* **108** (2002), no. 5-6, 733–754.
- [37] H.-K. Zhang, Current in periodic Lorentz gases with twists, *Comm. Math. Phys.* **306** (2011), no. 3, 747–776.

CEMAPRE, ISEG, UNIVERSIDADE TÉCNICA DE LISBOA, RUA DO QUELHAS
6, 1200-781 LISBOA, PORTUGAL

E-mail address: delmagno@iseg.utl.pt

DEPARTAMENTO DE MATEMÁTICA AND CEMAPRE, ISEG, UNIVERSIDADE
TÉCNICA DE LISBOA, RUA DO QUELHAS 6, 1200-781 LISBOA, PORTUGAL

E-mail address: jldias@iseg.utl.pt

DEPARTAMENTO DE MATEMÁTICA AND CMAF, FACULDADE DE CIÊNCIAS,
UNIVERSIDADE DE LISBOA, CAMPO GRANDE, EDIFÍCIO C6, PISO 2, 1749-016
LISBOA, PORTUGAL

E-mail address: pduarte@ptmat.fc.ul.pt

CEMAPRE, ISEG, UNIVERSIDADE TÉCNICA DE LISBOA, RUA DO QUELHAS
6, 1200-781 LISBOA, PORTUGAL

E-mail address: jpgaivao@iseg.utl.pt

CEMAPRE, ISEG, UNIVERSIDADE TÉCNICA DE LISBOA, RUA DO QUELHAS
6, 1200-781 LISBOA, PORTUGAL

E-mail address: dpinheiro@iseg.utl.pt